



Libraries and Learning Services

University of Auckland Research Repository, ResearchSpace

Version

This is the publisher's version. This version is defined in the NISO recommended practice RP-8-2008 <http://www.niso.org/publications/rp/>

Suggested Reference

Saeedi, H., Basher, Z., & Costello, M. J. (2016). Modelling present and future global distributions of razor clams (*Bivalvia: Solenidae*). *Helgoland Marine Research*, 70, 24 pages. doi: [10.1186/s10152-016-0477-4](https://doi.org/10.1186/s10152-016-0477-4)

Copyright

Items in ResearchSpace are protected by copyright, with all rights reserved, unless otherwise indicated. Previously published items are made available in accordance with the copyright policy of the publisher.

This is an open-access article distributed under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License.

For more information, see [General copyright](#), [Publisher copyright](#), [SHERPA/RoMEO](#).

ORIGINAL ARTICLE

Open Access



Modelling present and future global distributions of razor clams (Bivalvia: Solenidae)

Hanieh Saeedi^{*} , Zeenatul Basher and Mark J. Costello

Abstract

Razor clams (Pharidae and Solenidae) are deep-burrowing bivalves that inhabit shallow waters of the tropical, sub-tropical, and temperate seas. Using 'maximum entropy', a species distribution modelling software, we predicted the most suitable environments for the entire family and 14 *Solen* species to indicate their present and future geographic distributions. Distance to land, depth, and sea surface temperature (SST) were the most important environmental variables in training and creating the present and future distribution models both at the family and species level. In the present distribution models at the family level, the most suitable environment was where distance to land was between 0 and 100 km, a depth of 0–150 m, wave height of 5–7 m, a mean chlorophyll-*a* concentration about 0.7 mg m⁻³, and mean SST between 12 and 28 °C. Comparison with the future distribution models at the species level, found that most species were predicted to shift their distribution ranges poleward under the future environmental scenarios; i.e. species in the northern hemisphere would shift northward and southern species southward. Models also predicted that half of the species would expand their distribution ranges, 29% of species would not change their distribution, and 21% of species would shrink their distribution ranges under future climate change. Expanding geographic ranges would result in overlap in species ranges and thus greater species richness at regional scales. Model results predict that the mid-latitude peaks of species richness will move further apart, increasing the dip in richness near the equator, due to global climate change.

Keywords: Species distribution modelling, MaxEnt, Climate change, Range shifts, Mollusca, Ocean Biogeographic Information System, Global Biodiversity Information Facility

Background

Global climate change will influence the future distributions of marine species [1, 2]. Distribution maps indicating future suitable environments can predict the possible range shift of benthic species as a response to increasing temperatures [1, 3, 4]. Species are likely to respond to climate warming by shifting their distributions poleward [2, 5]. Predictive suitable environmental modelling is widely used to identify the environmental factors that control organisms' distribution and predict their geographic range from reported locations [3, 6–10].

One of the most ecologically and economically important superfamilies of marine Bivalvia is Solenoidea which

has two families, Pharidae and Solenidae, referred to as razor clams [11–13]. In an ecological context, razor clams' contributions to trophic food webs include serving as prey to crabs, gastropods, sea birds, and demersal fish [14, 15]. In some countries such as Iran, some species of razor clams (e.g., *Solen dactylus*) are being harvested by the local fishermen as bait for fishing and/or shrimp aquaculture [12, 16].

The family Solenidae has two genera, *Solen* Linnaeus, 1758 (65 accepted species) and *Solena* Mörch, 1853 (2 accepted species) with long narrow shells. Solenidae are deep-burrowing bivalves which dig to about 30 cm depth in low intertidal and subtidal sediments [11, 13, 16]. They have free-swimming larvae and may grow 20–30 mm per year [12]. Solenidae are mostly distributed in subtidal zones down to about 100 m in the tropical and temperate seas along the Indo-Pacific and Indian Ocean, east and north-west Pacific, European Atlantic coasts, western

*Correspondence: hanieh.saeedi@auckland.ac.nz; hanieh.saeedi@gmail.com
Institute of Marine Science, University of Auckland, Auckland 1142, New Zealand

south Atlantic Ocean, and north and south American coasts. They are absent from the polar regions in both the southern and northern hemispheres. The tropical Indo-Pacific area contains the highest number of species (about 75–80% of all known species) [11, 17].

Despite their wide distribution and ecological importance, limited sampling and ease of mis-identification (due to subtle morphological differences between species) contribute to gaps in knowledge of Solenidae global distribution patterns [15, 18, 19]. Environmental modelling may provide a better indication of the actual geographic distribution than reported locations alone, and can enable predictions of the effects of climate change.

Saeedi et al. (in press) found that the latitudinal gradient of species richness in razor clams was asymmetric and bimodal, with more species in the northern hemisphere and a dip between 0° and –15° latitude. Chaudhary et al. (2016) found this was typical for marine taxa, so razor clams may be a good model taxon for other marine species' biogeography [20]. Indeed, the biogeography of razor clams species' endemismity matched well that of marine species overall [21]. This study predicts the global distribution of the Solenidae family and 14 species based on environmental variables. We then test the hypothesis that species distributions would shift polewards from their present distributions under future climate change. Thus, our findings may be applicable to a wide range of marine taxa.

Methods

Geographic distribution

Data on species geographic distributions were gathered from the Global Biodiversity Information Facility (GBIF), Ocean Biogeographic Information System (OBIS), published literature, museum collections, and personal contacts. We cross-referenced OBIS and GBIF data to avoid duplication of records. We excluded all records that were classified as fossils, were mapped on land, had location coordinates that either had no precision estimates, or if location precision was more than 100 km. All species' names were verified in World Register of Marine Species (WoRMS), and their synonyms and misspellings were reconciled. We also re-examined razor clams in the collections of the Natural History Museum of Paris, Auckland Museum in New Zealand, the National Museum of Natural History (Smithsonian) in Washington, D.C., and the Natural History Museum of London. The specimens' identify was corrected if misidentified and geographic locations recorded. The museums were selected because of convenience of access (Auckland) and likelihood of holding large *Solen* collections. We found reliable coordinates for all 67 accepted *Solen* and *Solena* species that were listed in WoRMS. For the family level distribution

modelling, we also included *Solen gordonis* which is described as taxon inquirendum (a questionable species that requires revision and may or may not prove to be a valid species) in WoRMS. We further found an extra nine potential species described as aff. and cf. In total, 77 Solenidae taxa were used in this study [17].

Environmental data

Developing a species distribution model needs environmental variables which are likely to influence the species' distribution [22]. Environmental variables were selected regarding their relevance to Solenidae distribution and their biological importance in affecting *Solen* species populations and diversity [23]. As razor clams are distributed in coastal waters, distance to land, depth, salinity, pH, tidal height, and wave height could be important factors in limiting their distributions [14, 17, 24, 25]. While regionally depth and distance to land may be correlated, they are not globally ($r = -0.46$, $p \geq 0.05$, Additional file 1: Table S1) because of the variation in sizes of continental shelves and occurrence of mid-ocean islands. Sea surface temperature (SST), dissolved and saturated oxygen, and surface current may also affect the distribution, growth, reproduction, juvenile survival and mortality of *Solen* species [14, 17, 26–29]. Razor clams are filter feeders and consume phytoplankton as a food source. Thus, inorganic nutrients (such as silicate, nitrate, and phosphate), photosynthetically active radiation, diffuse attenuation coefficient (an indicator which shows how deeply visible light penetrates into the water column), chlorophyll-*a* concentrations, and primary productivity could indirectly or directly affect the distribution of Solenidae [14, 25, 30, 31]. Calcite was included because razor clams need calcium carbonate (CaCO_3) for shell growth [32].

Most environmental data layers were extracted from Global Marine Environment Datasets (GMED) [33] at a spatial resolution of 5 arcmin (0.083° grid cell pixel size, ca. 9 km at equator) (Table 1). Ocean area (km^2) and coastline length including islands (km) were extracted from Biogeoinformatics of Hexacorals (<http://www.kgs.ku.edu/Hexacoral/>) [34]. Layers were cropped to 70°N–70°S based on the maximum geographic distributions recorded for Solenidae species at latitudes 60°N and 50°S. Merow et al. (2013) found that, with the exception of SST range, using multiple derivatives of SST did not improve the performance of their model compared to using only a single derivation of SST [35]. For this reason, we calculated correlation coefficients of all selected GMED variables in ArcGIS using multivariate analysis (Band Collection Statistics) (Table S1), and select only one derivation of each metric, that is the mean, minimum, or maximum of temperature and chlorophyll-*a* concentration following Basher et al. (2014) [36]. Thus,

Table 1 List of environmental variables used in this study (from Basher et al. 2014) which were cropped to an extension of 70°N–70°S

Layer	Unit	Type	Temporal range	Minimum	Maximum	Mean	Std. Dev.
Land distance	km × 100	–	–	0.00	24.92	6.80	5.22
Sea surface temperature (annual mean)	°C	Monthly climatology	2002–2009	–1.00	31.54	15.89	10.44
Sea surface temperature (range)	°C	Monthly climatology	2002–2009	0.00	27.81	4.26	3.08
Depth	m	–	–	–10,293.65	0	–3671.68	1602.58
Wave height	m	–	–	0.00	7.00	0.29	1.02
Photosynthetically active radiation (annual mean)	Einstein m ⁻² day ⁻¹	Monthly climatology	1997–2009	0.00	64.82	35.22	8.55
Chlorophyll- <i>a</i> concentration (annual mean)	mg m ⁻³	Monthly climatology	2002–2009	0.02	0.90	0.05	0.04
Diffuse attenuation coefficient* (at 490 nm)	m ⁻¹	Monthly climatology	2002–2009	0.02	0.90	0.05	0.04
Primary Productivity	mg C m ⁻² day ⁻¹ cell ⁻¹	Annual climatology	–	0.00	4875.00	385.08	285.55
Tide average (average of maximum tidal height)	m	Annual climatology	–	0.00	6.38	0.51	0.44
Surface current	m s ⁻¹	Monthly climatology	2009–2010	–0.93	1.00	0.00	0.08
Salinity	PSS	In situ measure: 2009	1961–2009	0.00	41.00	33.96	2.09
pH	–	In situ measure: 2009	1910–2007	6.73	8.62	8.19	0.06
Dissolved oxygen	ml l ⁻¹	In situ measure: 2009	1898–2009	2.00	9.86	5.29	1.27
Saturated oxygen	ml l ⁻¹	In situ measure: 2009	1874–2000	76.05	113.11	100.10	3.23
Calcite concentration	mol m ⁻³	Seasonal climatology	2002–2009	0.00	9.00	2.87	3.18
Silicate	μmol l ⁻¹	In situ measure: 2009	1930–1986	0.00	69.00	9.85	13.86
Nitrate	μmol l ⁻¹	In situ measure: 2009	1922–1986	0.00	45.96	5.52	6.13
Phosphate	μmol l ⁻¹	In situ measure: 2009	1874–2000	0.00	2.00	0.26	0.44
<i>Sea surface temperature at 2100</i>	°C	Monthly climatology	2087 – 2096	0.00	35.05	18.04	10.91
<i>Salinity at 2100</i>	PSS	Monthly climatology	2087–2096	0.00	40.05	34.37	1.99
<i>Primary productivity at 2100</i>	mg C m ⁻² day ⁻¹ cell ⁻¹	Annual climatology	2090–2099	0.00	5004.00	354.76	277.07

Future environmental variables are in italics. Note that there were 19 variables for the Present, and 5 (including present land distance and depth as constant variables, and three future environmental variables in italics) for the Future model. Annual means were used and for sea surface temperature the annual range was also used as it was significantly different from the sea surface temperature annual mean. * An indicator of water clarity which expresses how deeply visible light in the blue to the green region of the spectrum penetrates into the water column. Distance to land was the distance to the nearest land using the Euclidean distance formula in ArcGIS

we only used annual mean SST and mean chlorophyll-*a* concentration.

In total, 19 environmental data layers were used to create the present day distribution models at both a family level and a species level (Table 1). However, for the future climate change scenarios there were only three environmental variables available, namely salinity, primary productivity, and SST mean. We assumed distance to land and depth would be similar until 2100, and thus predicted future geographic distributions by comparing the five variables including distance to land, depth, salinity, primary productivity, and SST mean for the present (reduced present day model) and future scenarios (Table 1). The differences between the mean values for the present and future environmental variables were 2 °C greater for sea surface temperature, 0.4 PSS less for salinity, and 30 mg C m⁻² day⁻¹ cell⁻¹ less for primary productivity (but 129 mg C m⁻² day⁻¹ cell⁻¹ greater maximum) (Table 1).

Modelling of species distributions

MaxEnt was selected in this study due to its success in developing species distribution models for marine species [37–40]. It has been widely used in conservation planning, ecology, evolution, epidemiology, invasive-species management and other fields [3, 6, 37]. MaxEnt minimizes the relative entropy, or dispersion, between two probability densities, one estimated from presence data, and one from the landscape in the context of covariate space. MaxEnt is optimized for predicting the realised or actual (rather than the fundamental) species distributions. Predictions of presence will thus still be dependent on the sample locations of the available data [7, 37]. Thus, any deficiency in sampling coverage might still bias the results [7, 41, 42].

MaxEnt version 3.3.3e was used to predict present and future (at year 2100) suitable environments for Solenidae on a global scale at both the family and species level. At the family level, a total of 526 distribution records of

77 Solenidae species were used for training the model [17] (see also Additional file 1: Table S2). Modelling at the family level allowed coverage of all species globally. Because of the greater number of distribution records the family level mapping would encompass each species level prediction. This if a species level model predicted a distribution outside the family level it would suggest poor model accuracy due to insufficient primary data.

MaxEnt was used to predict the suitable present and future environments for 14 *Solen* species which had more than 10 presence records separately (Table S2). We used one observation point per 0.083° pixel, to eliminate any duplicate points and reduce clumping. Models were created with 10 bootstrap replicates using default parameters for a random seed: randomly select 75% of the species presence records for training and 25% for testing the model in each replication stage [37, 43]. Then the average predictions across the all replicates were used for further analysis. The regularization multiplier was set to 1, and the maximum number of background points was increased to 100,000 instead of the default because of our large-scale mapping objective. There were 900 maximum iterations, and a convergence threshold of 0.00001 for the present day modelling [7, 23, 44]. We kept the default regularization values following Dudik et al. (2006) [45] as they result in better performance of evaluation data for presence only datasets. For the future projections, 10 cross-validated replicate models were generated. Default parameters including hinge features, random test percentage of zero [37, 43], and the other settings were the same as in the present day modelling.

To test the performance of MaxEnt models we used the Receiver Operating Characteristic (ROC) analysis. ROC analysis characterises the performance of a model at all possible thresholds using the Area Under the Curve (AUC) [6, 23, 44]. The highest numbers of AUC show more sensitive and specific model sets, ranging from 0.5 (random accuracy) to 1.0 (perfect discrimination) [7, 44]. We used the cumulative threshold value from the MaxEnt output which is a balance among training omission, predicted area, and threshold value. Values observed below the thresholds were considered to be unsuitable for the species. To determine the most important factors in training the distribution models and creating the final distribution models, we used the model outputs of the jackknife test as well as contribution rate (represents the importance of a given variable in model training), and permutation importance [46]. Permutation importance depends only on the final Maxent model (not for the replicates). The contribution for each factor is determined by randomly permuting the values of that factor among the points used for training the model and measuring the variation of AUC (training) value. A large decrease

indicates that the model depends heavily on that factor. Final values are normalized to a percentage for easier interpretation [46].

A map of environment suitability for Solenidae was generated to reflect the predicted probability of species occurrence using ArcGIS v10 [36, 43]. The model often predicted suitable environment in areas that were not contiguous with species' present distributions. For example, an Australian species in Japan, an Atlantic species on both coasts of north America, and Indian Ocean species in the Mediterranean. Such distributions are only likely if species are introduced by human activities. No marine species' are known to have larvae that disperse more than 1000 km [47, 48]. Thus, when determining both present and future predicted distribution ranges, only continuous distribution ranges within a maximum of 30° latitude (equivalent to 3300 km) and 30° longitude, and with more than a 75% prediction rate beyond the reported distribution were considered.

Results

Present model with 19 variables at family level

Distance to land and depth had the highest contribution rates and importance in creating the present distribution models at the family level (about 75%) (Additional file 1: Fig S1). After distance to land and depth, mean SST had the highest contribution rate (9.3%) in training the models, and wave height had the most importance rate in creating the final distribution models (Fig S1). The probability of Solenidae family occurrence in the present distribution models was close to 1 (the highest probability rate) where: distance to land was between 0 and 100 km; depth was 0–150 m; mean SST between 12 and 28 °C; wave height 5–7 m; and mean chlorophyll-*a* concentration 0.7 mg m⁻³ (Additional file 1: Fig S2).

Present model with 19 variables for 14 species

Species occurrences were strongly associated with wave height and distance to land which had the highest contribution rates in training the present distribution models of 10 species (about 70%) (Table 2). In the remaining four species, SST mean, SST range, dissolved oxygen, and depth had the highest contribution rates in training the present distribution models. Depth, distance to land, and calcite had the highest permutation importance in creating the final present distribution models (Table 2). The probability of *Solen* species occurrence in the present distribution models was close to 1 where: distance to land was approximately less than 200 km; depth less than 150 m; mean SST between 12 and 32 °C; primary productivity between 500 and 2500 mg C m⁻² day⁻¹ cell⁻¹, and salinity between 23 and 41 PSS (Table 3). In a total of 14 species, four cold temperate species including *Solen*

Table 2 The output of highest contribution and importance values of environmental variables in creating the present MaxEnt distribution models for 14 species using 19 variables

Solen species	Training records	Contribution		Permutation importance	
		Highest	Rate (%)	Highest	Rate (%)
<i>aureomaculatus</i>	15	Wave height	22.8	Depth	32.2
<i>canaliculatus</i>	12	SST mean	19.0	Land distance	39.2
<i>foresii</i>	50	Wave height	22.2	Calcite	28.7
<i>grandis</i>	24	SST range	53.2	Land distance	56.7
<i>kajiyamai</i>	12	Dissolved oxygen	29.6	Depth	56.1
<i>marginatus</i>	126	Land distance	42.8	Calcite	34.5
<i>roseomaculatus</i>	35	Wave height	33.6	Land distance	84.7
<i>sicarius</i>	30	Land distance	60.9	Depth	67.4
<i>sloanii</i>	15	Land distance	60.9	Depth	67.4
<i>strictus</i>	17	Land distance	28.0	Depth	46.4
<i>thuelchus</i>	12	Wave height	25.1	Land distance	52.0
<i>vagina</i>	10	Land distance	28.4	Calcite	21.6
<i>vaginoides</i>	85	Depth	24.4	Calcite	38.5
<i>viridis</i>	33	Wave height	29.3	Land distance	33.2

Values are normalized to give percentages. The average AUC of training data was from 0.97 to 1 from the ten model runs, with little variation between runs indicating a good model fit

grandis, *S. marginatus*, *S. thuelchus*, and *S. viridis* preferred the lowest temperature ranges from 12 to 16 °C, two warm temperate species including *Solen sicarius* and *S. strictus* had a temperature preferences from 16 to 19 °C, and the rest were tropical species with high temperature preferences from 25 to 32 °C. Four species of 14 species including *Solen canaliculatus*, *S. roseomaculatus*, *S. sloanii*, and *S. vagina* favoured high salinities from 40 to 41 PSS, *S. sicarius* preferred the low salinity from 23 to 25 PSS, and the rest of species mostly predicted in salinities from 29 to 39 PSS (Table 3).

Reduced present models with 5 variables for 14 species

Depth and distance to land had the highest contribution rates in training and creating the reduced present distribution models for almost all *Solen* species, except for *Solen marginatus* where mean SST had the highest importance permutation rate in creating the final distribution models (Table 4).

Future models with 5 variables at family level

Distance to land and depth had the highest contribution rates and importance in creating the future distribution models at the family level (about 85%) (Fig S1). Mean SST had the highest contribution rate (11.7%) in training the future projections after distance to land and depth. However, primary productivity was the third important factor in creating the future distribution models after distance to land and depth (Fig S1). The probability of Solenidae family occurrence in the future distribution models was

close to 1 where: distance to land was between 0 and 100 km; depth 0–150 m; mean SST 12–28 °C; primary productivity between 500 and 2000 mg C m⁻² day⁻¹ cell⁻¹; and salinity between 30 and 38 PSS (Additional file 1: Fig S3).

Future models with 5 variables for 14 species

In all 14 species excluding *S. aureomaculatus*, distance to land had the highest contribution rate in training the future distribution models (Table 4). However, mean SST was the most important factor in creating the final future distribution models in half of the 14 *Solen* species (50%). After mean SST, depth in five species (36%), and distance to land in two species (14%) were the most important factors in creating the final future distribution models. The probability of *Solen* species future distributions was close to 1 where distance to land was less than 200 km; depth less than 150 m; mean SST 11–29 °C; primary productivity between 500 and 2300 mg C m⁻² day⁻¹ cell⁻¹; and salinity between 32 and 42 PSS (Table 3).

The maximum averages of predicted suitable primary productivity for *Solen canaliculatus*, *S. kajiyamai*, *S. marginatus*, and *S. roseomaculatus* were lower in the future distribution models compared to the present distribution models (Table 3). Predicted suitable salinities in the future distribution models were higher compared to the present distribution models in all species, except for *S. canaliculatus*. The maximum predicted suitable SST mean in half of the species was lower in the future distribution models compared to the present distribution

Table 3 The variables that had the highest predicted probability of *Solen* species occurrence for the present and future distribution models

<i>Solen</i> Species	Depth (m)	Present			Depth (m)	Future		
		Primary productivity (mg C m ⁻² day ⁻¹ cell ⁻¹)	Salinity (PSS)	Mean sea surface temperature (°C)		Primary productivity (mg C m ⁻² day ⁻¹ cell ⁻¹)	Salinity (PSS)	Mean sea surface temperature (°C)
<i>aureomaculatus</i>	0–150	500–750	35–36	26–28	0–150	500–1000	40–42	25–28
<i>canaliculatus</i>	0–150	2000–2500	40–41	29–31	0–150	750–1000	32–34	25–28
<i>foresii</i>	0–150	800–1100	34–35	27–30	0–150	800–1100	35–36	25–29
<i>grandis</i>	0–150	1900–2000	30–32	13–14	0–150	1800–2000	30–33	13–14
<i>kajiyamai</i>	0–150	1800–2000	39–40	30–31	0–150	1500–1700	40–41	22–25
<i>marginatus</i>	0–100	1500–1900	39–40	12–13	0–100	1400–1600	40–41	11–13
<i>roseomaculatus</i>	0–150	1400–1600	40–41	31–32	0–150	1000–1200	41–42	23–26
<i>sicarius</i>	0–150	2200–2300	23–25	16–17	0–150	2200–2300	32–33	15–17
<i>sloanii</i>	0–100	750–850	40–41	25–26	0–100	750–1000	41–42	23–25
<i>strictus</i>	0–100	1800–2000	29–31	17–19	0–100	1500–1700	32–33	18–19
<i>thuelchus</i>	0–100	2200–2100	39–41	13–16	0–100	2200–2000	40–41	15–16
<i>vagina</i>	0–150	1600–1900	40–41	25–28	0–150	1600–2100	41–42	26–28
<i>vaginoides</i>	0–100	800–1100	38–40	26–28	0–100	800–1100	39–41	26–27
<i>viridis</i>	0–100	2000–2100	38–39	12–14	0–100	2000–2200	38–40	12–13

Distance to land was approximately less than 200 km for all species in both present and future distribution models

models, and it did not change for the remainder. Temperature species such as *Solen grandis*, *S. marginatus*, *S. strictus*, *S. sicarius*, *S. thuelchus*, and *S. viridis* had the lowest predicted suitable SST ranges from 12 to 19 °C in both present and future distribution models (Table 3).

Present distributions

More than 50% of species showed similar present predicted distribution ranges compared to their actual distribution ranges (Fig. 1). MaxEnt distribution model outputs at the family level indicated that the most suitable environments for Solenidae at present are in the shallow waters of the northern east Pacific (California, USA), tropical west Atlantic (north Carolina, USA), European Atlantic, Gulf of Thailand, and eastern (Brisbane and Melbourne) and western coast of Australia (Additional file 1: Fig S4).

Future distributions

Solen sloanii and *S. roseomaculatus* were distributed in both hemispheres and had the widest latitudinal distribution ranges of all the species (Figs. 1, 2). They were predicted to shift northward in the northern, and southward in the southern hemisphere under the future climate change scenarios (Table S2, Figs. 1, 2, 3). *Solen vagina* seems to be an endemic species to the Gulf of Thailand and was predicted to shift southward to Malaysia, Singapore, and Indonesia. *Solen strictus* and *S. grandis* showed


similar future distribution ranges compared to their current distributions. *Solen canaliculatus* was predicted to shift northward from its present distributions and disappear from Taiwan in the future distribution model. High latitude species such as *Solen marginatus*, *S. sicarius*, and *S. viridis* were predicted to shift northward in the northern, and *S. thuelchus* southward in the southern hemisphere. However, the extent of their predicted distribution ranges in the present and future distribution models were similar. All Australian species including *Solen foresii*, *S. kajiyamai*, *S. aureomaculatus*, and *S. vaginoides* were predicted to shift southward along the eastern and southern Australia coasts in the future models such that the species' would split into east and west coast populations. All Australian species disappeared from the northern part of Australia under future climate change (Table S2, Figs. 1, 2).

At the family level, more geographic areas were predicted to be suitable for razor clams in the future compared to the present distribution models (Additional file 1: Fig S5). This is supported by the results of individual species models. Considering predicted present and future distribution models at the species level, half of the 14 species had a wider distribution range in the future compared to predicted present distribution models (Fig. 3). In contrast, three species (21%), namely *Solen vaginoides*, *S. strictus*, and *S. viridis* were predicted to have narrower latitudinal distribution ranges in the

Table 4 Output contribution and importance values of five environmental variables used to create the reduced present day and future MaxEnt distribution models for 14 *Solen* species

Solen species	Training records	Present					Future				
		Environmental variables Contribution rate/permutation importance (%)					Environmental variables Contribution rate/permutation importance (%)				
		Depth (m)	Land distance (km x 100)	Primary productivity (mg C m ⁻² day ⁻¹ cell ⁻¹)	Salinity (PSS)	Mean sea surface temperature (°C)	Depth (m)	Land distance (km x 100)	Primary productivity (mg C m ⁻² day ⁻¹ cell ⁻¹)	Salinity (PSS)	Mean sea surface temperature (°C)
<i>aureomaculatus</i>	13	53/16	21/78	8/2	0.2/0.0	18/5	35/68	35/3	7/3	2/0.3	22/25
<i>canaliculatus</i>	10	23/86	52/1	4/0.1	4/0.9	17/12	13/1	56/28	4/2	5/7	21/62
<i>foresii</i>	41	54/11	33/86	0.0/0.0	0.0/0.0	12/3	30/56	46/7	4/1	3/3	16/32
<i>grandis</i>	20	22/8	37/83	0.6/0.1	1.6/2	25/6	27/47	39/13	12/7	7/5	15/29
<i>kajiyamai</i>	10	64/58	18/39	3/0.2	0.4/0.0	14/2	25/47	53/12	4/2	0.8/0.4	17/39
<i>marginatus</i>	105	21/4	44/8	14/1	7/3	14/84	2/2	62/29	12/2	5/3	19/65
<i>roseomaculatus</i>	11	52/8	33/90	0.0/0.0	0.7/0.0	14/2	18/10	62/41	0.2/0.4	0.2/0.2	19/49
<i>sicarius</i>	24	4/0.1	56/89	23/2	2/1	14/8	3/1	58/57	23/11	3/6	14/25
<i>sloanii</i>	12	42/3	46/95	0.7/0.0	0.6/0.0	11/2	7/0.6	67/59	2/0.5	1/0.2	24/40
<i>strictus</i>	12	58/21	36/78	0.1/0.0	0.4/0.1	5/1	11/12	59/37	2/0.4	9/8	18/42
<i>thuelchus</i>	10	56/17	38/83	0.1/0.0	1/0.2	4/0.2	6/5	65/38	7/3	0.0/0.1	21/54
<i>vagina</i>	6	49/15	25/80	0.0/0.0	2/0.2	24/5	11/3	63/37	0.4/0.0	4/2	22/58
<i>vaginoides</i>	69	16/90	54/2	7/1	6/2	16/5	21/69	49/8	8/3	5/5	17/15
<i>viridis</i>	28	10/84	53/8	11/0.2	6/0.6	20/8	18/5	47/30	9/0.8	5/4	20/60

Values are normalized to give percentages. Environmental variables with highest contribution and permutation importance rates represent the most important variables in training and creating the final distribution models. The average AUC of training data in both present and future distribution models was from 0.97 to 1 from the ten model runs, with little variation between runs indicating a good model fit



future. Four species (29%) had similar present and future distribution ranges indicating that their distribution would not change due to future climate change. A species range may change but the mean latitude of its ranges may or may not change. The mean latitudes of the northern hemisphere species were predicted to change showing a negligible change in the future (Fig. 4). However, the mean latitude of southern species would shift southward.

Discussion

Present distributions

The majority of the predicted suitable environments were in the shallow waters of the temperate and tropical north America, Indo-West Pacific, and the European Atlantic Ocean where Solenidae have been reported. Some of these areas may not be inhabited by *Solen* at present due to dispersal constraints. For example, there were no distribution records of Solenidae in New Zealand; although the model predicted this area had a suitable environment now and in the future. This would

suggest that this family had not occurred on Gondwanaland or evolved prior to the separation of New Zealand from Australia (around 80 million years ago). During glaciations, SST in northern New Zealand were above 10 °C [49] which is within the temperature tolerance of Solenidae. Evidently, the duration and/or behaviour of Solenidae planktonic larvae might have been insufficient for species to colonise New Zealand from Australia.

The environment suitability model developed in this study indicated that Solenidae species' distributions were highly correlated with distance to land, depth, SST, and wave height. Distance to land and depth contributed to over 70% of the variation in the models. Mean SST had the next most contribution rate (about 9%) to the global models after distance to land and depth. SST is a major factor in the reproduction, larval development, recruitment, and mortality of Solenidae [14, 31].

The models predicted Solenidae to occur in coastal areas (≤ 200 km from the land) with depths of less than




Fig. 2 The latitudinal distribution ranges of *Solen* species according to field data (continuous lines), the predicted present (dashed lines) and future distribution (dotted lines) range with more than 75% prediction rates. The central line shows the Equator. The distribution median (open circle) and mean (triangle and cross) are indicated





Fig. 3 The predicted present and future latitudinal ranges of 14 *Solen* species. Symbols above the line indicate a wider latitudinal range, and symbols below the line indicate a predicted decrease in the species latitudinal range. Black circles: southern latitudes (5 species); open circles: northern latitudes (9 species)

150 m, wave height of 5–7 m, mean SST of 12–28 °C, and primary productivity of 400–700 mg C m⁻² day⁻¹ cell⁻¹). The suitable environments for Solenidae species were similar to the temperate razor clam *Ensis directus* (Phariidae) [50]. They also found high probabilities of occurrence for *E. directus* at depths between 0 and 67 m, minimum annual SSTs between 3 and 18 °C and maximum annual SSTs between 20 and 26.5 °C [50].

Wave height had the highest contribution rate in training the present distribution models of most high latitude species including *Solen aureomaculatus*, *S. fonesii*, *S. thuelchus*, and *S. viridis*. The greater occurrence of Solenidae in areas with 5–7 m wave height, which is at the upper end of the range of potential wave height, may be an indicator of the occurrence of sandier sediments. Nickerson (1975) reported that densities of razor clams were highest on sandy beaches with least silt, and that silt-laden sediments might be responsible for suffocation of razor clams in early life stages [51].

Future distributions

Mean SST was the most important environmental variable in half of the species in creating the final future distribution models. The future species distribution models showed that northern and southern hemisphere species would shift northward and southward respectively under future climate warming. Half of the species were predicted to expand their distribution ranges 21% of species to shrink, and 29% of species did not change their distribution under future climate change. The potential poleward range shifts due to global warming have been also reported for sandy-beach invertebrates [5] and tropical molluscs [2]. Indo-West Pacific areas (especially China Sea, Sea of Japan, Bay of Bengal, Gulf of Thailand, Andaman Sea, Philippines, Indonesia, and Papua New Guinea)



would be occupied by more *Solen* species due to the warming average temperatures, as predicted for other tropical molluscs [2]. Distributions of tropical *S. canaliculatus*, *S. vagina*, *S. strictus*, and *S. grandis* could also expand northward and southward along the coastal areas of the China Sea, Sea of Japan, Bay of Bengal, and Gulf of Thailand under future climate change. However, physical geographical barriers would limit future distribution changes into these areas [11, 18].

Almost all Australian species were predicted to shift their distribution southwards and disappear from its northern territory. The sea surface temperature in northern Australia is predicted to become 2–3 °C warmer (33 °C) in the future [52] which would be out of these species temperature tolerance [17]. In contrast, the predicted future distribution models of some high latitude species, such as *S. marginatus* and *S. sicarius*, would not expand outside their current distribution ranges.

The present and future distribution models in *Solen roseomaculatus* and *S. sloanii* showed exceptionally wide disjunct distributions. These merit confirmation because they may reflect misidentifications as a consequence of their subtle differences in shell morphology and lack of molecular taxonomy studies. Thus they could be more than one species [17, 53].

Conclusions

Saeedi et al. (in press) found that the latitudinal distribution of Solenidae was bimodal, with most species at the edges of the tropics. They suggested this may be typical for marine species in general, because such bimodality has also been found for taxa as varied as planktonic

foraminifera and marine mammals [20]. The results of the present study show that this bimodality is likely to increase due to climate warming, and will result in increased species richness at regional scales because most species will increase their geographic range. Thus climate warming can be considered as an unintended global experiment that confirms the role of temperature in defining the latitudinal distribution of marine species.

Additional file

Additional file 1. Input environmental variables and geographic distribution data, and output results of the present and future distribution modelling. This file includes the additional data on correlations between the environmental variables, input geographic distribution points of Soleinidae species used in the present and future modelling, and the models' output which could not be fitted inside the paper due the page limits.

Abbreviations

MaxEnt: maximum entropy; SST: sea surface temperature; GBIF: Global Biodiversity Information Facility; OBIS: Ocean Biogeographic Information System; WoRMS: World Register of Marine Species; GMED: Global Marine Environment Datasets; PSS: practical salinity scale; ROC: receiver operating characteristic; AUC: area under the ROC curve.

Authors' contributions

HS: study design, data collection, statistical analysis, species distribution modelling, writing the paper. ZB: environmental data interpretation, species distribution modelling, review of the paper. MJC: study design, statistical analysis, writing of the paper. All authors read and approved the final manuscript.

Acknowledgements

HS was supported by the New Zealand International Doctoral Scholarship (NZIDRS) and University of Auckland Doctoral Scholarship. We would like to thanks the referees for their helpful comments.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The dataset supporting the conclusions of this article is included within the article (and its Additional file 1).

Funding

This research was funded by the New Zealand International Doctoral Scholarship (NZIDRS) and University of Auckland Doctoral Scholarship. The funding bodies had no roles in the design of the study and collection, analysis, interpretation of data, and in writing the manuscript.

Received: 20 May 2016 Accepted: 4 November 2016

Published online: 20 December 2016

References

- Cheung WWL, Lam VWY, Sarmiento JL, Kearney K, Watson R, Pauly D. Projecting global marine biodiversity impacts under climate change scenarios. *Fish Fish.* 2009;10(3):235–51.
- Saupe EE, Hendricks JR, Peterson AT, Lieberman BS. Climate change and marine molluscs of the western North Atlantic: future prospects and perils. *J Biogeogr.* 2014;41(7):1352–66.
- Rombouts I, Beaugrand G, Dauvin JC. Potential changes in benthic macrofaunal distributions from the English Channel simulated under climate change scenarios. *Estuar Coast Shelf Sci.* 2012;99:153–61.

4. Sarà G, Milanese M, Prusina I, Sarà A, Angel DL, Glamuzina B, Nitzan T, Freeman S, Rinaldi A, Palmeri V, Montalto V, Lo Martire M, Gianguzza P, Arizza V, Lo Brutto S, De Pirro M, Helmuth B, Murray J, De Cantis S, Williams GA. The impact of climate change on mediterranean intertidal communities: losses in coastal ecosystem integrity and services. *Reg Environ Change*. 2014;14(Suppl 1):5–17.
5. Schoeman DS, Schlacher TA, Defeo O. Climate-change impacts on sandy-beach biota: crossing a line in the sand. *Glob Change Biol*. 2014;20(8):2383–92.
6. Phillips SJ, Dud M, Schapire RE. A maximum entropy approach to species distribution modeling. Proceedings of the twenty-first international conference on Machine learning; Banff, Alberta, Canada. 1015412: ACM; 2004. p. 83.
7. Phillips SJ, Anderson RP, Schapire RE. Maximum entropy modeling of species geographic distributions. *Ecol Model*. 2006;190(3–4):231–59.
8. Elith J, Leathwick JR. Species distribution models: ecological explanation and prediction across space and time. *Annu Rev Ecol Syst*. 2009;40(1):677–97.
9. Ashcroft MB, French KO, Chisholm LA. An evaluation of environmental factors affecting species distributions. *Ecol Model*. 2011;222(3):524–31.
10. Davies AJ, Guinotte JM. Global habitat suitability for framework-forming cold-water corals. *PLoS ONE*. 2011;6(4):e18483.
11. Cosel Rv, editor. An introduction to the razor shells (Bivalvia: Solenacea). Hong Kong: Hong Kong University Press; 1990.
12. Saeedi H, Raad SP, Ardalan AA, Kamrani E, Kiabi BH. Growth and reproduction of *Solen dactylus* (Bivalvia: Solenidae) on northern coast of the Persian Gulf (Iran). *J Mar Biol Assoc UK*. 2009;89(8):1635–42.
13. Saeedi H, Costello MJ, Cosel Rv. First report of anterior pallial tentacles in *Solen dactylus* (Bivalvia: Solenidae) from the Northern Persian Gulf, Iran. *PLoS ONE*. 2013;8(5):e63487.
14. Guerra A, Lodeiros C, Gaspar MB, da Costa F. Razor clams: biology, aquaculture and fisheries. Xunta de Galicia, Consellería do Mar, Santiago de Compostela. Spain 2011.
15. Lassuy DR, Newell RIE, Fritzsche RA, Hassler TJ, Simons D, Group USAEW-ESCE, et al. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest): Pacific razor clam: Defense Technical Information Center; 1989.
16. Saeedi H, Ardalan AA, Kiabi HB, Zibaseresht R. Metal concentrations in razor clam *Solen dactylus* (Von Cosel, 1989) (Bivalvia: Solenidae), sediments and water in Golshahr coast of Bandar Abbas, Persian Gulf. *Iran J Fish Sci*. 2012;11(1):165–83.
17. Saeedi H, Costello MJ, Dennis T. Bimodal latitudinal species richness and high endemism in razor clams (Bivalvia: Solenidae). *J Biogeogr. (to appear)*.
18. Cosel Rv. Seven new species of *Solen* (Bivalvia: Solenidae) from the tropical Western Pacific, with remarks on other species. In: 301–341; col. pls. 31–38. In: Falkner MG, K. & Speight, C.D., editor: *Collectanea Malacologica - Festschrift für Gerhard Falkner: 10 + 547 pp + 48 col. Pls.* Conchbooks, Hackenheim/Frierich-Held-Gesellschaft, München; 2002.
19. Saeedi H, Costello MJ. Aspects of global distribution of six marine bivalve mollusc families. In: da Costa F, editor. *Clam fisheries and aquaculture*. New York: Nova Science Publishers Inc.; 2013. p. 27–44.
20. Chaudhary C, Saeedi H, Costello MJ. Bimodality of latitudinal gradients in marine species richness. *Trends Ecol Evol*. 2016;31(9):670–6.
21. Costello MJ, Tsai P, Wong PS, Cheung A, Basher Z. Marine biogeographic realms and species endemism. *BioScience. (to appear)*.
22. Ready J, Kaschner K, South AB, Eastwood PD, Rees T, Rius J, Agbayani E, Kullander S, Froese R. Predicting the distributions of marine organisms at the global scale. *Ecol Model*. 2010;221(3):467–78.
23. Byers JE, McDowell WG, Dodd SR, Haynie RS, Pintor LM, Wilde SB. Climate and pH predict the potential range of the invasive apple snail (*Pomacea insularum*) in the Southeastern United States. *PLoS ONE*. 2013;8(2):e56812.
24. Lewin J, Eckman JE, Ware GN. Blooms of surf-zone diatoms along the coast of the Olympic Peninsula, Washington. XI. Regeneration of ammonium in the surf environment by the Pacific razor clam *Siliqua patula*. *Mar Biol*. 1979;52(1):1–9.
25. da Costa F, Martínez-Patiño D. Culture potential of the razor clam *Solen marginatus* (Pennant, 1777). *Aquaculture*. 2009;288(1–2):57–64.
26. Remacha-Triviño A, Anadón N. Reproductive cycle of the razor clam *Solen marginatus* (Pulteney 1799) in Spain: a comparative study in three different locations. *J Shellfish Res*. 2006;25(3):869–76.
27. Abrahão JR, Cardoso RS, Yokoyama LQ, Amaral ACZ. Population biology and secondary production of the stout razor clam *Tagelus plebeius* (Bivalvia, Solecurtidae) on a sandflat in southeastern Brazil. *Zoologia (Curitiba)*. 2010;27:54–64.
28. Saeedi H, Kamrani E, Matsuoka K. Catastrophic impact of red tides of *Cochlodinium polykrikoides* on razor clam *Solen dactylus* in coastal waters of northern Persian Gulf. *J Persian Gulf (Mar Sci)*. 2011;2(6):13–20.
29. da Costa F, Nóvoa S, Ojea J, Martínez-Patiño D. Changes in biochemical and fatty acid composition of the razor clam *Solen marginatus* (Solenidae: Bivalvia) during larval development. *Mar Biol*. 2011;158(8):1829–40.
30. Trainer VL, Cochlan WP, Erickson A, Bill BD, Cox FH, Borchert JA, Lefebvre KA. Recent domoic acid closures of shellfish harvest areas in Washington State inland waterways. *Harmful Algae*. 2007;6(3):449–59.
31. Hmida L, Ayache N, Haouas Z, Romdhane MS. Oocyte cohort analysis: criteria for an evaluation of the reproductive cycle in *Solen marginatus* (Pennant, 1777), (Bivalvia: Solenacea) in Southern Tunisia. *J Shellfish Res*. 2010;29(1):129–34.
32. Lomovskiy BJ, Gutiérrez JL, Iribarne OO. Identifying repaired shell damage and abnormal calcification in the stout razor clam *Tagelus plebeius* as a tool to investigate its ecological interactions. *J Sea Res*. 2005;54(2):163–75.
33. Basher Z, Costello MJ, Bowden DA. Global marine environment dataset (GMED). World Wide Web electronic publication. Version 1.0 (Rev.01.2014). <http://gmed.auckland.ac.nz>. Accessed 2014-05-12.
34. Fautin DG, Malarky L, Soberón J. Latitudinal diversity of sea anemones (Cnidaria: Actiniaria). *Biol Bull*. 2013;224(2):89–98.
35. Merow C, Smith MJ, Silander JA. A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography*. 2013;36(10):1058–69.
36. Basher Z, Bowden DA, Costello MJ. Diversity and distribution of deep-sea shrimps in the Ross sea region of Antarctica. *PLoS ONE*. 2014;9(7):e103195.
37. Phillips SJ, Dudík M. Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography*. 2008;31(2):161–75.
38. Cordellier M, Pfenninger M. Inferring the past to predict the future: climate modelling predictions and phylogeography for the freshwater gastropod *Radix balthica* (Pulmonata, Basommatophora). *Mol Ecol*. 2009;18(3):534–44.
39. Pineda E, Lobo JM. The performance of range maps and species distribution models representing the geographic variation of species richness at different resolutions. *Glob Ecol Biogeogr*. 2012;21(9):935–44.
40. Basher Z, Costello MJ. The past, present and future distribution of a deep-sea shrimp in the Southern Ocean. *PeerJ*. 2016;4:e1713.
41. Franklin J. Mapping species distributions spatial inference and prediction/ Janet Franklin with contributions by Jennifer A. Miller. In: Miller JA (ed) Cambridge: Cambridge University Press; 2009.
42. Slater H, Michael E. Predicting the current and future potential distributions of *Lymphatic filariasis* in Africa using Maximum Entropy ecological niche modelling. *PLoS ONE*. 2012;7(2):e32202.
43. Elsässer B, Fariñas-Franco JM, Wilson CD, Kregting L, Roberts D. Identifying optimal sites for natural recovery and restoration of impacted biogenic habitats in a special area of conservation using hydrodynamic and habitat suitability modelling. *J Sea Res*. 2013;77:11–21.
44. Oliveira MH, Calheiros SK, Jacobi CM, Latini RO. Modeling the potential distribution of the invasive golden mussel *Limnoperna fortunei* in the Upper Paraguay River system using limnological variables. *Braz J Biol*. 2010;70(3):831–40.
45. Dudík M, Schaapire SJ, Phillips SJ. Correcting sample selection bias in maximum entropy density estimation. *Adv Neural Inf Process Syst*. 2006;18:323–30.
46. Phillips SJ. A brief tutorial on Maxent, versions: 3.3.1. <http://www.cs.princeton.edu/~schapire/maxent/on>. Accessed 2016-07-26.
47. Kinlan BP, Gaines SD. Propagule dispersal in marine and terrestrial environments: a community perspective. *Ecology*. 2003;84(8):2007–20.
48. Gaines SD, Gaylord B, Gerber LR, Hastings A, Kinlan B. Connecting places: the ecological consequences of dispersal in the sea. *Oceanography*. 2007;20(3):90–9.

49. Barrows TT, Juggins S. Sea-surface temperatures around the Australasian margin and Indian Ocean during the Last Glacial Maximum. *Quat Sci Rev*. 2005;24:1017–47.
50. Raybaud V, Beaugrand G, Dewarumez JM, Luczak C. Climate-induced range shifts of the American jackknife clam *Ensis directus* in Europe. *Biol Invasions*. 2015;17(2):725–41.
51. Nickerson RB. A critical analysis of some razor clam (*Siliqua patula* Dixon) populations in Alaska. Juneau: Alaska Department of Fish and Game; 1975.
52. Johnson JE, Holbrook NJ. Adaptation of Australia's marine ecosystems to climate change: using science to inform conservation management. *Int J Ecol*. 2014;2014:1–12.
53. Rv Cosel. Three new species of *Solen* (Bivalvia: Solenidae) from the Indian Ocean, with remarks on the Solenidae of Madagascar. *J Conchol*. 1989;33(4):189–208.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit



Table S1. Correlation coefficients between the environmental variables. SST: Sea Surface Temperature; PAR: Photosynthetically Active Radiation; DA: Diffuse Attenuation Coefficient; Chl-*a*: chlorophyll-*a* concentration.

Environmental Variable	Salinity	Silicate	SST (mean)	SST (range)	Tide Average	Surface Current	Wave Height	Nitrate	Dissolved Oxygen	Saturated Oxygen	PAR (mean)	pH	Phosphate	Primary Productivity	Depth	Land Distance	DA (mean)	Chl- <i>a</i> (mean)	
Silicate	-0.24																		
SST (mean)	0.36	-0.74																	
SST (range)	-0.28	-0.24	-0.04																
Tide Average	-0.03	0.23	-0.03	-0.06															
Surface Current	-0.08	0.14	-0.31	0.04	-0.11														
Wave Height	-0.17	-0.09	0.01	0.29	0.12	-0.02													
Nitrate	-0.21	0.87	-0.86	-0.25	0.05	0.27	-0.13												
Dissolved Oxygen	-0.41	0.73	-0.96	0.04	0.04	0.30	0.00	0.82											
Saturated Oxygen	0.12	-0.80	0.54	0.35	-0.21	-0.05	0.10	-0.69	-0.51										
PAR (mean)	0.32	-0.63	0.92	-0.12	-0.00	-0.42	0.03	-0.77	-0.89	0.46									
pH	0.20	-0.42	0.42	0.10	0.01	-0.10	-0.03	-0.47	-0.41	0.36	0.35								
Phosphate	-0.21	0.78	-0.77	-0.28	0.02	0.28	-0.12	0.88	0.74	-0.62	-0.68	-0.44							
Primary Productivity	-0.29	-0.15	0.04	0.42	0.16	-0.02	0.51	-0.17	-0.02	0.18	0.06	-0.08	-0.17						
Depth	-0.26	0.09	-0.16	0.19	0.26	-0.00	0.47	0.02	0.18	-0.09	-0.09	-0.09	0.01	0.46					
Land Distance	0.13	-0.06	-0.08	-0.12	-0.27	0.13	-0.33	0.12	0.03	0.01	-0.12	0.00	0.09	-0.33	-0.46				
DA (mean)	-0.53	0.20	-0.30	0.37	0.17	0.01	0.44	0.13	0.32	-0.11	-0.23	-0.25	0.09	0.68	0.50	-0.33			
Chl- <i>a</i> (mean)	-0.53	0.20	-0.30	0.37	0.17	0.01	0.44	0.13	0.32	-0.11	-0.23	-0.25	0.09	0.68	0.50	-0.33	1		
Calcite	0.33	-0.45	0.67	-0.12	-0.07	-0.24	-0.16	-0.57	-0.65	0.33	0.65	0.34	-0.49	-0.29	-0.27	0.06	-0.43	-0.43	

Table S2. Geographic distributions of 14 *Solen* species based on field data and predicted present and future (in the year of 2100) environment suitability models. The last two columns column show only predicted suitable environments of the species which had the highest prediction rates ($\geq 75\%$ prediction rate) in MaxEnt. Locations predicted beyond their recorded distribution are bolded.

Species	Number records	Recorded distribution	Predicted present	Predicted future
<i>Solen aureomaculatus</i> Habe, 1964	19	Australia [Northern Territory: Gulf of Carpentaria, Groote Eylandt; Western Australia: Dampier Archipelago (East Lewis Island, off Enderby Island, South-West of Rocky Head, Dolphin Island, near East end of Searipple Passage, large bay on East coast, Georgeff Reef); Queensland: Townsville (Rowes Bay, Magnetic Is; Torres Strait, Great Barrier Reef, Cairncross Is, off Cape York Peninsula, Black beach); New South Wales: Kima]	Australia [Northern Territory; Western Australia; Queensland [Brisbane to Townsville]; New South Wales]	Australia [Western Australia [Perth]; Queensland; New South Wales [Sydney]; South Australia [Adelaide]]
<i>Solen canaliculatus</i> Tchang & Hwang, 1964	15	China [South China [Hong Kong and Hainan]; Taiwan [SW Tungkang]	China; Taiwan	China [Hangzhou]
<i>Solen fonesii</i> Dunker, 1862	60	Australia [Queensland: Dingo Bay; Broom; Brisbane; Hervey Bay; Keppel Bay; Townsville; Family Ids, National Park, Dunk Is; Cape York, Telegraph Station; GBR, Whitsunday Passage, Lindeman Is; Armstrongs Beach, Sarina; Lucinda; Seaforth; Town beach, Mackay, Boyne Island; Cannonvale Beach; Emu Creek, Sinclair Bay; Magnetic Island, Horseshoe Bay; Cardwell, Kennedy National Park; Brampton Reef, Bowen; Western Australia: Port Headland; Eighty (Ninety) Mile Beach, Wallal Downs; Buccaneer Archipelago; Onslow area; Carnarvon; SE of Exmouth Homestead; Northern Territory: Gulf of Carpentaria; Darwin; New South Wales: Gloucester Is; Black beach, Kima]	Australia [Western Australia; Queensland; Northern Territory]	Australia [Queensland; Western Australia [Perth]]

<i>Solen grandis</i> Dunker, 1862	30	Japan [Mikawa; Sagami; Harima; Katase-Enoshima, Fujisawa, Kanagawa Prefecture; Ishikawa, Honshu Island; Aichi, Mikawa; Wakayama, Nada-Cho]; China [Laoshan Bay, Qingdao coast; Intertidal Zone Kimje; Shandong Peninsula, coastal regions of the Yellow Sea; Rizhao, Shandong Province; Beihai City, Beibu Gulf of Guangxi, Southern China]; Republic of Korea [Buan-gun, Jeollabuk-do]	Japan; China; Republic of Korea	Japan; China; Republic of Korea
<i>Solen kajiyamai</i> Habe, 1964	15	Australia [Western Australia: Onslow Jetty, Onslow; Dolphin Island; North section of East coast; North of Nelson Rocks, Dampier Archipelago; Queensland: Dingo beach; Cape York "township"; North Keppel Is, off NW end Keppel Bay; Yeppoon, Keppel Bay; Turkey beach, between Bundaberg and Gladstone; Karumba; beaches around Mapoon; Sweers Is, Gulf of Carpentaria; Northern Territory]	Australia [Western Australia; Queensland; Northern Territory]	Australia [Western Australia [Shark Bay ; Perth]; Queensland; South Australia [Adelaide]; New South Wales [Sydney]]
<i>Solen marginatus</i> Pulteney, 1799	151	France [Arcachon Bay; Marennes-Oléron; Oyster-farming bay; Corse, Golfe de Porto, Vecchio; Berck]; Spain [Atlantic coast of Huelva; Barra de Cangas; Pontevedra, Boiro; Ría de Arousa; Regueira; Asturias; Vila Real de Santo Antonio; Galicia; Ria de Rebadeo; Santander Bay; Cantabria; Sotavento; Melilla; A Pobra do Caraminal, A Coruna]; Egypt [Port Said]; UK [Chichester Harbour; Porth Dinllaen; Lleyn Peninsula; Milford Haven, Pembrokeshire; Tenby, Pembrokeshire; Hampshire] ;Tunisia [12 k N Raoued; Gammarth; Maltin Wadi in Sfax]; Belgium; Morocco [Bou Regreg estuary; Mellila]; Italy [Adria bei Monfalcone; Istrien; Angle Bay; Ferrara; Porto Garibaldi; Forli; Cesenatico; La Eo ria; Rimini]; Turkey [Kocaeli; Izmit Korfezi]; Cyprus; Greece [Kavalla; Thasos; Alyki, Zakynthos; Laganas; Saronikos; Saronikos]; Netherlands; Belgium; Ireland; Portugal	France; Spain; Egypt; UK; Tunisia; Belgium; Morocco; Italy; Turkey; Cyprus; Greece; Netherlands; Belgium; Ireland; Portugal	France; Spain; Egypt; UK; Tunisia; Belgium; Morocco; Italy; Turkey; Cyprus; Greece; Netherlands; Belgium; Ireland; Portugal; Norway

[Averio; Faro; Ria Formosa lagoon]

<i>Solen roseomaculatus</i> Pilsbry, 1901	17 Yemen [Abyan beach, Aden]; Madagascar [Tuléar; 35 mi W Cap St Andro; 25 mi NW of Cap St Andre]; Mozambique [Inhaca, Portuguese Island; 80 mi SSW Ponta da Barra; At Linga-Linga Morrumbene in channel N of wrecked schooner]; Iran [Bandargah, Kish, The Persian Gulf]; Japan [Takehara, Hiroshima; Hirado, Hizen; Okinawa; Nagasaki]; Indonesia [Off West Coast of wasir Island, W Wocam, Aru]; Philippines [Marin Bay, Mindoro]; Ceram [Off Tutuhuhur, Piru Bay]	Yemen; Madagascar; Mozambique; Iran; Japan; Indonesia; Philippines; Ceram	Yemen; Madagascar; South Africa ; Mozambique; Iran; United Arab Emirates ; Japan; China
<i>Solen sicarius</i> Gould, 1850	36 United States [California (Long Beach; Santa Barbara; Monterey Bay; Cabrillo Beach); Washington (Olympia; Puget Sound; Victoria)]; Canada [Naden Harbor; Graham Island; Virago Sound; Tasu Sound; W of Moresby Island; Sandspit Bar, Deadtree Point; Naden Harbour; Skidegate]	United States [Washington; Baja California]; Canada [Vancouver]	United States; Canada
<i>Solen sloanii</i> Gray, 1843	19 Japan [Ryukyu Islands; Akashi Harima; Amamura, Awaji; 1km WNW Onna Vilage (Horseshoe Cliffs), Okinawa]; Mozambique [80 mi SSW Ponta da Barra; Inhaca]; Madagascar [Tuléar; Tanjon' Andavaka; Taolagnaro; Marolinta]; South Africa [Kosi Bay off; SE of Sheffield Beach; SE of Umhlanga Beach; Cape St. Lucia off]; Indonesia [Lembeh-Strait, NE Sulawesi]; Vietnam [Nha Trang]	Japan; Madagascar; South Africa; Indonesia; Vietnam	Japan [Tokyo]; South Africa [Port Elizabeth]; Indonesia; Vietnam
<i>Solen strictus</i> Gould, 1861	19 Japan [Enoshima; Chiba ; Nagasaki; Hakodate; Honmoku, Hirado, Hizen; Tokyo Bay, Honshu; Takasago; Yamaguchi, Yamaguchi Prefecture; Nago City, Okinawa Prefecture; TSU city, Mie prefecture beach; Takou; Aomori; Aichi, Mikawa]; China [Cheffo; Qingdao (Tsingtao)]; Malaysia [Sarawak]; Korea [Kushima, Ariake Bay (North Pacific Ocean); Cheju-do; Simpo	Japan; Malaysia; South Korea	Japan; Malaysia; South Korea

		tideland, Outfal Estuary of Kumkang River, West of Korea]		
<i>Solen thuelchus</i> Hanley, 1842	15	Brazil [Codo Beach; Rasa da Cotinga Island; Santana's Archipelago; Rio Grande do Sul; Praia do Camaroeiro; Praia Pereque]; Argentina [Mar Del Plata, Northern Argentina; Desembocadura Río Negro]; Uruguay [Costa de Maldonado]	Brazil [State of Rio de Janeiro to State of Santa Catarina; Macapá]; Argentina; Uruguay	Brazil; Argentina; Uruguay
<i>Solen vagina</i> <i>Linnaeus, 1758</i>	10	Malaysia [Kuantan estuary, Tanjung Lumpur, Mangrove area, Phang; George Town, Penang]; India [Calcutta; Digha Coast]; Vietnam [Nha Trang; Chu Lei Bay, Da Nang]; Singapore [Changi beach park, Changi]; Philippines [Negros Occidental, Cadiz]; Thailand [Andaman Sea, Krabi, Ban Laem]; Myanmar (Burma) [Maungmagan; Tavoy]; Andaman & Nicobar Islands	Malaysia [Brunei]; India; Vietnam; Singapore; Philippines; Thailand; Myanmar; Andaman & Nicobar Islands	Malaysia [Brunei]; India [South India]; Sri Lanka ; Vietnam; Singapore; Philippines; Thailand; Myanmar; Andaman & Nicobar Islands
<i>Solen vaginoides</i> Lamarck, 1818	104	Australia [Queensland: Brisbane; Keppel Bay; Townsville; Great Barrier Reef ; Wonga Beach, S end, N of Mossman; Sandy Strait State School, Urangan; Caloundra; offshore of Sandy Point, S of Mooan, Fraser Is; Tannum Sands, S of Gladstone; Pialba; Shelly Beach, Torquay, Hervey Bay; Port Curtis, off Gatcombe Head; Lucinda; Cape Flattery; Torres Strait, Friday Is; Bowen; Disaster Bay; Dingo beach; Telegraph Station; Somerset Bay, Cape York; Kurrimine Beach, S of Innisfail; Mission Beach, near Tully; Flat Rock, North Richmond River entrance; Rudder Reef 30 miles NE of Mossman; Amity point, N. stradbroke Is; Long Beach, Joskeleigh; New South Wales: Sydney; Victoria; Botany Bay; Twofold Bay; Jervis Bay; Eden; Newtons Beach, Nadgee Fauna Reserve; Merimbula, off Surfshed; Off Nelson Bay, Port Stephens; In Great Oyster Bay; Seven Mile Beach, W side of Frederick Henry Bay; Mallacoota; Bundeena beach; Gloucester Is; Broulee, S of Batemans Bay; Eastern Bass Strait, 3.2 km S of	Australia [Queensland; New South Wales; Southern Australia; Northern Territory; Western Australia [Perth]]	Australia [Queensland; New South Wales; Southern Australia; Western Australia]

Cape Conran; St Francis Island, Nuyts Arch; Cave Bay, Hunter Island, off NW tip; 13.3 km E of Eastern edge of Lake Tyers; 15.3 km ESE of Eastern edge of Lake Tyers; 20.2 km WSW of Pt Ricardo; 5.4 km SW of Cape Conran, Eastern Bass Strait; Crib Point, Western Port, CPBS station 32N; Between Stony Point and Crib Point; Off Cheseapeake, French Island; Chinamans beach; Port Phillip Bay, Sorrento pier; Tuncurry, Wallis Lake; Southern Australia: Tasmania; Adelaide; Melbourne; Northern Territory: (beaches around Mapoon; Groote Eylandt, Gulf of Carpentaria); Western Australia: Shoalwater Bay, off Canoe Passage between Townshend & Marquis Ids; Swan Is, W side, Shoalwater; Duke Group, Marble Is, Homestead Bay; Albany

Solen viridis
Say, 1821

40 United States [Sullivan's Bay, South Carolina; Newport, Rhode Island; Galveston Island, Texas; Smith Island; Holden Beach, New Carolina; Simone Island; Newport, Rhode Island; Georgia; Delaware Bay; Charleston bay, South Carolina; Beaufort Harbour, North Carolina; Atlantic City; Off 5th Saint & Beach, Wildwood; South Cape May beach; Anglesea; Egg Harbor, New Jersey; Hog I, Philadelphia; Jekyll Island; near Turner's Creek, Wilmington River; one mile offshore of S end of Ossabaw Isl, Georgia; Bay Marchand Lease Area, 500-2000 Ms N Of Platform]; Canada

United States [Carolina, Gulf of Mexico] ; Canada

United States [Eastern coasts]; Canada [**Nova Scotia**]





Fig. S1 Jackknife test of environmental variable importance and their contribution to predicted global Solenidae distribution at present (top) and in the future (bottom). Estimations of the relative contribution of each variable to the global model are given in front of the each bar. White bars show the performance of the global model (test gain) without each variable and black bars show the influence of each variable in isolation. Values shown are averages over 10 replicate runs. Mean sea surface temperatures (SST) show the highest contribution rates in creating both present and future distribution models, after depth and distance to land. Apart from distance to land and depth, wave height and primary productivity had the highest importance rates in creating the present and future distribution models, respectively.

Probability of Presence






Fig. S2 Marginal response curves of the present predicted probability of Solenidae occurrence, for explanatory environmental variables that contributed substantially to the global maximum entropy. The value shown on the y-axis of the response curves (between 0 to 1) is predicted probability of suitable conditions. The highest predicted probability values in each response curve show the most suitable environmental conditions.

Probability of Presence




Fig. S3 Marginal response curves of the future predicted probability of Solenidae occurrence, for explanatory environmental variables that contributed substantially to the global maximum entropy.




Fig. S4 Present predicted distribution of 77 Solenidae species extracted from Saedi et al. (in press) (probability of occurrences = 0 - 0.93). This family level map contains the distribution records of all species which even had only one distribution record.




Fig. S5 Future predicted distribution of 77 Solenidae species extracted from Saeedi et al. (in press) (probability of occurrences = 0 - 0.77). This family level map contains the distribution records of all species which even had only one distribution record.