



Libraries and Learning Services

# University of Auckland Research Repository, ResearchSpace

## Version

This is the Accepted Manuscript version. This version is defined in the NISO recommended practice RP-8-2008 <http://www.niso.org/publications/rp/>

## Suggested Reference

Wills, P. R. (2016). A Hilly path through the thermodynamics and statistical mechanics of protein solutions. *Biophysical Reviews*, 8(4), 291-298.  
doi: [10.1007/s12551-016-0226-6](https://doi.org/10.1007/s12551-016-0226-6)

## Copyright

The final publication is available at Springer via  
<http://link.springer.com/article/10.1007%2Fs12551-016-0226-6>

Items in ResearchSpace are protected by copyright, with all rights reserved, unless otherwise indicated. Previously published items are made available in accordance with the copyright policy of the publisher.

For more information, see [General copyright](#), [Publisher copyright](#), [SHERPA/RoMEO](#).

A Hilly path through the thermodynamics and statistical mechanics of protein solutions.

Peter R Wills

Department of Physics, University of Auckland, PB 92109, Auckland 1142,  
Aotearoa-New Zealand

[p.wills@auckland.ac.nz](mailto:p.wills@auckland.ac.nz)

[tel +64 9 9238889](tel:+6499238889)

[fax +64 9 3737445](tel:+6493737445)

## Abstract

The opus of Don Winzor in the fields of biophysical and analytical biochemistry is a major component of that certain antipodean approach to this broad area of research that blossomed in the second half of the twentieth century. The need to formulate problems in terms of thermodynamic nonideality posed the challenge of describing a clear route from molecular interactions to the parameters that biochemists routinely measure. That required delving into the statistical mechanics of solutions of macromolecules and at every turn mathematically complex, rigorous, general results that had previously been derived previously, often by Terrell Hill, came to the fore. Central to this work were the definition of the “thermodynamic activity”, the pivotal position of the polynomial expansion of the osmotic pressure in terms of molar concentration and the relationship of virial coefficients to details of the forces between limited-size groups of interacting molecules. This was richly exploited in the task of taking account of excluded volume and electrostatic interactions, especially in the use of sedimentation equilibrium to determine values of constants for molecular association reactions. Such an approach has proved relevant to the study of molecular interactions generally, even those between the main macromolecular solute and components of the solvent, by using techniques such as exclusion and affinity chromatography as well as light scattering.

## Key words

nonideality; thermodynamic activity; osmotic virial coefficient; excluded volume; sedimentation equilibrium; molecular association reaction

## Historical Prolegomena

When I arrived at the John Curtin School of Medical Research at the Australian National University at the end of 1977, armed with nothing more than a PhD thesis about protein hydrodynamics and dynamic light scattering (DLS), along with a few equations describing a polymerizing system at thermodynamic and sedimentation equilibrium, the name of Sandy Ogston was still writ large over the portal of the Department of Physical Biochemistry and the incumbent Head, Laurie Nichol, was engraving his own tablets of stone with the laws of excluded volume and thermodynamic nonideality. But I soon discovered that another fellow called Don Winzor regularly came down from the University of

Queensland at Brisbane to act as the immovable stone on which Laurie sharpened his intellectual chisels. The two sat together for hours going over pages of manuscripts and arguing about a multitude of thermodynamic expressions and their meaning, always trying to tease out some new experimental consequence of the eternally intractable relation between strictly defined theoretical quantities and their messy measureable counterparts. When I showed some propensity for mathematical reasoning I was invited to enter the inner sanctum and was admitted into the cabal of those initiated in the rites of “thermodynamic intuition”. The three of us ended up writing a joint paper about non-sigmoidal Scatchard plots (Nichol, Wills, Winzor, 1979), but the main task I took on was to develop a more complete account of thermodynamic non-ideality in terms of more accurately estimated activity coefficients of separate species in interacting systems, a study that brought McMillan-Mayer theory (2) into focus and eventually gave what seemed like quite a good description of lysozyme’s self-association in mildly acidic solutions (Wills, Nichol, Siezen, 1980). After a couple of years I took off to a new job in Berlin, but having acquired the new skill of making linear and quadratic order corrections to the standard ideal-solution equations I penned a note with Nichol and Ogston about the effect of inert polymers on protein self-association (Nichol, Ogston, Wills, 1981) and applied the technique to DLS data (Wills, Georgalis, 1981) all with a view to rationalizing the mostly impossible behavior of aggregation-prone ribosomal proteins. However, the completion of this last task had to wait another decade and a half, that is, until Don was prepared to slice through the experimental messiness of it all and convince us that the results really did mean something (Wills, Georgalis, Dijk, Winzor, 1995). In the meantime, the science of excluded volumes and their effect on nucleic acid melting was meted out in the dark, smoky atmosphere of a West Berlin *Kneipe* known as The Flying Dutchman, reminiscent of a locale in a Le Carré novel, conveniently located on the Richard-Wagner-Strasse in Charlottenburg, just over from the Deutsche Oper. The obliging barman Roland spirited documents back and forth between authors and journal editor, notifying us whenever a new package with mysteriously arcane contents was ready for pick-up (Woolley, Wills 1985). Such were the ups and downs of life for a young biochemist in those days, looking for meaning in activity coefficients without thinking hard enough to realize that there was more than one to be had ...

And so it came to pass that when Laurie Nichol left science to become a university administrator his parting gift to me was to attach Don Winzor to my hip and convince us that we should (and could) find a proper solution to the problem of unifying the not-completely-rigorous biochemical description of macromolecular interactions in terms of excluded volume, electrostatic and molecular association effects, unaware as we were that the task had long been completed and we had occasionally used bits of the relevant Hill papers without comprehending their full relevance. But first Don had an old score to settle. We would show that one could be protected from all of the complications of “preferential solvation” by staying under the aegis of excluded volume (Winzor, Wills, 1986). And by then I had joined the molecular biological heretics who were interested in prions (Wills, 1986) and strayed off on sabbatical to Carleton Gajdusek’s lab at the National Institutes of Health in Maryland. Those were days of glorious intellectual freedom when a scientist paid to study neurological

disease could live a parallel life, not only incurring the wrath of the Pentagon and eventually the FBI by using the Freedom of Information Act to play amateur WikiLeaks, but also pursuing thermodynamic rigour, strolling over in the next building to say hullo to Allen Minton only to be introduced to the legendary Terrell Hill. We now had a wise and patient guide to show us the way, someone who had blazed the trail a quarter century earlier. It took him no time to explain our naïvety in thinking that the expression for the thermodynamic activity of a single component could be transferred willy-nilly across to multicomponent systems. No, one had to decide how the system was constrained while the concentration of a component was varied. Nowhere was this more simply evident than in the definition of the osmotic pressure (Figure 1). Armed with this new perspective and the help of Wayne Comper, who already knew how to do proper thermodynamics using differential quantities, we could finally reconcile measurements of virial coefficients made under alternative “osmotic” or isobaric conditions (Wills, Comper, Winzor, 1993). We could then describe preferential solvation and the effect of small molecules on macromolecules rigorously (Wills, Winzor, 1993) and this in turn took us back to the basic theory of sedimentation equilibrium and the problem how the concentration dependence of the buoyancy term  $(1 - \bar{v}\rho)$  was fudged during integration of  $d\ln c/dr^2$  to get the equation of the radial concentration dependence  $c(r)$  of a non-ideal solute. After years of claim and counterclaim we could finally resolve that as long as one stuck to the molar concentration scale, the equation for a non-ideal solution could be obtained from the ideal equation simply by replacing  $c(r)$  with the osmotic activity  $z(r)$ , at least in the case of an incompressible solution, and then the density appearing in the buoyancy factor is unequivocally that of the pure solvent, rendering the buoyancy term independent of concentration. And then we really were in a position to entrench the antipodean perspective that non-ideality could not be dispensed with by curve-fitting (Adams, Fujita, 1963), but had to be dealt with prior to any realistic assessment of the extent of molecular association reactions. A strong collaboration was underway.

## Thermodynamic activity of a solute

The ideal equation for the chemical potential  $\mu_A$  of a solute  $A$

$$\mu_A = \mu_A^o + RT \ln X_A \quad (1)$$

is not particularly sensitive to the choice of the scale on which the concentration  $X$  is nominally measured, simply because the most commonly used scales (amount of solute per amount of solution, or alternatively, per amount of solvent) converge within a linear correction as  $X \rightarrow 0$  and use of Eq. (1) to describe the thermodynamic behaviour of a solute depends on the extent of the dilute regime in which corrections that are linear in  $X$  can be neglected. For example, as long as the specific volumes of both solute  $v_A$  and solvent  $v_s$  are effectively constant under the relevant experimental conditions, the molar concentration  $C$  (or its mass per unit volume equivalent  $c_A = C_A M_A$ ) are related to the molal concentration  $m_A$  through the equation

$$C_A = m_A \rho_s / (1 + M_A v_A \rho_s m_A) \quad (2)$$

where  $\rho_s = 1/v_s$  is the density of the solvent and  $M_A$  is the molar mass of the solute. Eq. (1) can be modified to reflect the non-ideal effects of molecular interactions simply by replacing the nominal concentration variable with a thermodynamic “activity”, which is usually considered to be the product of the concentration and an “activity coefficient”, a fudge factor that corrects the concentration to give the correct value of the chemical potential. However, to obtain a quantitatively useful expression involving a thermodynamic activity it is necessary to decide how the solution is to be constrained as the solute concentration is varied, a choice that corresponds to determining the standard state to which  $\mu^\circ$  refers (see Fig. 1). It turns out that the two most useful choices for the standard state of the solute are the infinite dilution limit under isothermal conditions at some specified value of either the chemical potential of solvent  $\mu_s$  or the pressure  $P$ . In the first case the natural concentration scale on which to define the “osmotic activity”  $z_A = \gamma_A C_A$  is the molar scale (moles of solute per unit volume of solution)

$$\mu_A(T, \mu_s, C_A) = \mu_A^\circ(T, \mu_s) + RT \ln z_A \quad (3)$$

and in the second case the natural concentration scale on which to define the “isobaric activity”  $a_A = \gamma_A m_A$  is the molal scale (moles of solute per unit mass of solvent)

$$\mu_A(T, P, m_A) = \mu_A^\circ(T, P) + RT \ln a_A \quad (4)$$

because in each case Eq. (1) is preserved as the ideal expression for  $\mu_A$ , at least for an incompressible solution ( $v_A$  and  $v_s$  constant), applicable to the constraint (cont.  $\mu_s$  or  $P$ ) under which the concentration is varied.

The statistical mechanics of solutions of macromolecules under osmotic and isobaric conditions was given a thorough treatment by Hill in the late 1950s (Hill, 1956a; 1958; 1959). He provided a very general analysis, showing how the coefficients,  $B_2$ ,  $B_3$ , etc. appearing in the virial expansion for the osmotic pressure, that is, changes in  $P$  due to changes in  $C_A$  at constant  $\mu_s$ ,

$$[\Pi/(RT)]_{T, \mu_s} = C_A (1 + B_2 C_A + B_3 C_A^2 + \dots) \quad (5)$$

can be related to the coefficients  $C_2$ ,  $C_3$ , etc. in the equivalent expansion that accounts for changes in  $\mu_s$  due to changes in  $m_A$  at constant  $P$ ,

$$-[(\mu_s - \mu_s^\circ)/(\rho_s RT)]_{T, P} = m_A (1 + C_2 m_A + C_3 m_A^2 + \dots) \quad (6)$$

The great advantage of the osmotic pressure expansion, Eq (5), is that any virial coefficient  $B_n$  can, in principle, be calculated directly by consideration of the

energy of interaction between just  $n$  molecules of solute. However, either expansion can be used to define the activity coefficient appropriate to conditions under which the corresponding activity is defined:

$$\ln \gamma_A = 2B_2C_A + (3/2)B_3C_A^2 + \dots \quad (7)$$

and

$$\ln y_A = 2C_2m_A + (3/2)C_3m_A^2 + \dots \quad (8)$$

whence a molecular interpretation can eventually be made of variations in the thermodynamic activity of a macromolecular solvent.

These results, and their application to a wide variety of problems, were by no means the exclusive domain of Hill. Indeed, he stood on the shoulders of the likes of Scatchard (1946), Brinkman and Hermans (1949), Bird, Spotz and Hirschfelder (1950), Stockmayer (1950) and Kirkwood and Buff (1951), to name but a few. However, the most systematic and comprehensive development of McMillan-Mayer theory in relation to protein solutions was carried out by Hill (1954; 1955a;b; 1956a;b; 1957; 1958; 1959), the main results of relevance to biochemists being incorporated into his specialized book (Hill, 1968) and further elaborated later (Hill, Chen, 1973).

### Sedimentation Equilibrium

With this clarification of what was meant by the term “thermodynamic activity” it was possible to take a new approach to the derivation of the expression for the concentration distribution of a solute in a system at sedimentation equilibrium, giving a clear comparison with the standard result for an ideal solution:

$$C_A(r) = C_A(r_0)\psi(r) \quad (9a)$$

where

$$\psi(r) = \exp[\phi(r^2 - r_0^2)] \quad (9b)$$

and

$$\phi = M_A(1 - v_A\rho_s)\omega^2/(2RT) \quad (9c)$$

The two usual ways of obtaining Eq (9), the sedimentation-diffusion flux balance method and the thermodynamic equilibrium method, both involve an integration step in which the solution density  $\rho(r)$  is treated as a constant, independent of  $r$ , when it is not. For incompressible solutions,

$$\rho(r) = \rho_s + (1 - v_A\rho_s)M_AC_A. \quad (10)$$

The mathematical sleight of hand involved is of little consequence in the analysis of data that can be considered to represent the behaviour of an ideal solution and it accounts for versions of Eq (9) in which  $\rho$  is said variously to be the solvent or the solution density. However, in deriving an expression for the exact

magnitude of the first order nonideality correction to the sedimentation equilibrium concentration profile, the dependence of  $\rho$  on  $C_A$  must be taken into account correctly. The exact result for incompressible solutions can be written simply by replacing  $C_A$  in Eq (9) with the osmotic activity  $z_A$ :

$$z_A(r) = z_A(r_0)\psi(r) \quad (11)$$

We found a very general pathway to this result (Wills, Jacobsen, Winzor, 1996), and devised a direct method for the analysis of sedimentation equilibrium data based on a transformation to  $\psi(r)$  from Eq (9b) as the independent variable. We also presented a derivation for the simplest case of single solute (Wills, Hall, Winzor, 2000), as well as extending the  $\psi(r)$  analysis to mixtures of interacting solutes (Wills, Jacobsen, Winzor, 2000).

To understand the relevance of the osmotic activity in sedimentation equilibrium it is important to realise that the standard chemical potential  $\mu_A^\circ(T, \mu_s)$  in Eq (3) is a function of radial distance  $r$ . Consider a solution of the same composition and at the same temperature and pressure as that existing at  $r_0$ . Remove all of the solute under conditions of constant chemical potential of solvent  $\mu_s(r)$ , to give pure solvent at a pressure of  $P^0(r_0) = P(r_0) - \Pi(r_0)$ . According to Eq (3) this osmotic change has brought the solute to its standard state of infinite dilution, where its chemical potential is  $\mu_A^\circ(T, \mu_s)$  as at  $r_0$  in the centrifuge cell. Carry out the same procedure for a solution corresponding to the conditions at  $r$ . Now, the hypothetical change  $\Delta\mu_A^\circ(T, \mu_s)$  in the chemical potential of *solute* between these two states of the solvent is  $RT\phi(r^2 - r_0^2)$ , with the density in Eq (9c)

unequivocally  $\rho_s$  because the change of state corresponds to the ideal situation in which a single solute molecule in an infinite volume of solvent is transformed from the solvent conditions of  $P^0$  and  $\mu_s$  at  $r_0$  to those at  $r$ . Eq (11) follows immediately from the calculation of  $\mu_A(r) - \mu_A(r_0)$  for the conditions of the actual sedimentation equilibrium under consideration.

## Molecular interactions

For molecules that interact through a spherically symmetric potential  $u(x)$  when separated by a distance  $x$ , the molar-scale second virial coefficient ( $L$  is Avogadro's number) is given by (McMillan, Mayer, 1945)

$$B_2 = -2\pi L \int_0^\infty f(x)x^2 dx \quad (12)$$

where the Mayer  $f$ -function is defined as

$$f(x) = 1 - \exp[-u(x)/kT] \quad (13)$$

and for the hard-sphere interaction between molecules of radius  $R_A$

$$u(x) = \begin{cases} \infty & x < 2R_A \\ 0 & x \geq 2R_A \end{cases} \quad (14)$$

the van der Waals excluded volume per pair of molecules

$$B_2 = 16\pi LR_A^3/3. \quad (15)$$

is obtained (Wills, Winzor, 2005).

The contribution of excluded volume to thermodynamic non-ideality had long been understood and Ogston and Winzor (1975) had extended Eq. (15) to the case of ellipsoids and used Debye-Hückel theory to take account of charge-charge interactions. However, in the background were much earlier discussions in which Hill (1954; 1956b) had already provided a simple way of calculating the third virial coefficient  $B_3$  as well as a more realistic estimate of charge-charge effects that could easily be adapted to macromolecular solutes. The form of the electrostatic potential adopted for molecules with a net surface charge of  $Z_A$  multiplied by the electronic charge  $e$  was

$$u(x) = \frac{Z_A^2 e^2 \exp[-\kappa(x - 2R_A)]}{\epsilon x (1 + \kappa R_A)^2} \quad x \geq 2R_A \quad (16)$$

where  $\kappa$  is the Debye-Hückel inverse screening length of the supporting electrolyte-bearing medium and  $\epsilon$  its dielectric constant. This was demonstrated to give a good representation of the dependence of  $B_2$  on ionic strength (Wills, Hall, Winzor, 2000) and provided confidence in what had become a common practice of using the relevant formulae to “calculate out” the effects of excluded volume and charge, thereby bringing into visibility molecular associations due to short range attractive forces between proteins. Comparison of different ways of taking into account higher order effects of charge-charge interactions, through either extensions of Eq (16) (Hill, 1954; 1956b) or scaled particle theory emphasised the need for independent information about the size and charge of proteins in attempting to investigate association reactions (Scott, Wills, Winzor, 2010).

Once we had rediscovered the Hill-Chen opus (Hill, Chen, 1973) and its theoretical underpinnings, Laurie Nichol’s “unification” problem was solved. Molecular associations could indeed be treated as just another form of nonideality, exactly in the manner envisaged by the van der Waals equation of state (Wills, Winzor, 2005). In the case of a single solute we were not required to have separate thermodynamic equations for different oligomeric “components” as had been done previously (Wills, Nichol, Siezen, 1980); all we had to do was incorporate the dimerization constant  $K_2$  into the relevant virial coefficients

$$B_2 = B_{11}^* - K_2; \quad B_3 = B_{111}^* - 2K_2(4B_{11}^* - B_{12}^*) + 4K_2^2; \quad \text{etc.} \quad (17)$$



and concomitantly more complex relationships for higher order coefficients. The star superscript in Eq (17) denotes a virial coefficient that is calculated solely on the basis of the relatively long range repulsive forces (excluded volume and electrostatic) between molecules according to the likes of Eqs (15) & (16); and the string of subscripts on these quantities indicate a list of monomers (1), dimers (2), etc. comprising a total of  $n$  monomer units.

With this basic insight we were able to extend the direct analysis of sedimentation equilibrium data using the  $\psi$ -function of Eq (11) as far as the case of multiple experiments involving two separate components (ovalbumin and cytochrome *c*) under conditions in which they undergo an association reaction to form a hetero-dimer (Wills, Jacobsen, Winzor, 2000). We also gave some consideration to  $B_3$  and higher order effects (Wills, Winzor, 2001; Wills, Scott, Winzor, 2012). The  $\psi$ -function analysis predated other procedures for the direct calculation of association constants through statistical analysis of sedimentation equilibrium data, emphasising the non-thermodynamic nature of the separation of the short-range molecular forces involved in molecular association reactions from others (Winzor, Wills, 2007); and thereby invalidating statistical curve fitting as a way of eliminating effects due to other interactions as a way of access to equilibrium constants. The direct use of Eq (17) to extract protein dimerization constants from osmotic pressure data was also demonstrated (Wills, Winzor, 2009).

### Other extensions

The standard analysis (Cassasa, Eisenberg, 1964) of the thermodynamic effects exerted by the electrolyte routinely regarded as part of the solvent in which a protein is dissolved was legendary among biochemists for its rigour and complexity. However, our analysis of “preferential solvation” (Winzor, Wills, 1986; Wills, Winzor, 1993) and its application to sedimentation analysis (Jacobsen, Wills, Winzor, 1996) had led us to an approach using molar-scale variables and osmotic considerations (Wills, Winzor, 2002) than was much simpler than the corresponding approach based on the use of molal quantities. Provided the components of the diffusible osmotic solvent could be regarded as a simple volume-filling ideal mixture, their only significant effect was to alter the effective density of the medium in which the macromolecules were immersed, giving rise to an altered solvent density

$$\rho_d = \rho_s + (1 - v_M \rho_s) M_M C_M \quad (18)$$

and an effective protein specific volume  $v_A^*$ , defined in terms of an altered sedimentation buoyancy factor, which depended on the 2<sup>nd</sup> virial coefficient  $B_{AM}$  for the interaction between molecules of *A* and *M*

$$M_A (1 - v_A^* \rho_d) = M_A (1 - v_A \rho_s) - (1 - v_M \rho_s) B_{AM} M_M C_M + \dots \quad (19)$$

This approach, which was found to be in good agreement with experiment, is vindicated by the interpretation of Eq (4.37) of Hill (1968). In the case of sedimentation equilibrium the derivation of Eq (11) is still valid for a multicomponent solvent, provided one applies the osmotic condition of constant chemical potential to all such components. When they are all small enough not to undergo significant sedimentation then the buoyancy factor is a constant throughout the cell. The presence of larger, inert polymers added to mimic the crowded molecular environment inside a cell could be handled in a similar, very simple way, provided they made the overwhelmingly dominant contribution to the non-ideality of the protein (Wills, Georgalis, Dijk, Winzor, 1995; Winzor, Wills 2006).

Thermodynamic results are general, so insights in the context of thinking about what happens in a centrifuge cell can be transferred to an array of other experimental situations. The partitioning of a solute in a frontal chromatography experiment is clearly reminiscent of osmosis and lends direct access to the effects of nonideality. This had been exploited in the analysis of “preferential solvation” (Winzor, Wills, 1986) and the use of inert polymers in the estimation of the size of proteins (Wills, Georgalis, Dijk, Winzor, 1995; Winzor, Wills, 2006), but it was also applied in a careful analysis of haemoglobin self-association at high concentrations (Winzor, Wills, 2003) and extended to the new technique of self-interaction chromatography (Winzor, Scott, Wills, 2007), once again proving how consideration of excluded volume could provide a parsimonious explanation of diverse phenomena, including the influence of an inert polymer on the kinetics of enzymic catalysis, an effect that had previously been ascribed to “osmotic stress” (Winzor, Wills, 1995). Failure to take proper account of the definition of the thermodynamic activity, osmotic or isobaric, Eqs (3) and (4), and the concentration scale against which virial coefficients are measured, molar or molal, Eq (2), remains an impediment to the optimal interpretation of thermodynamic measurements on protein solutions (Wills, Winzor 2011). The interpretation of data from light scattering experiments (Winzor, Deszczynski, Harding, Wills, 2007) remains incomplete.

## **Conclusion**

The contribution of Don Winzor to more than half a century of physical and analytical biochemistry, starting from considerations of excluded volume and continuing to demand that statistical mechanical expositions lead to experimentally relevant conclusions and interpretations, has been indispensable to the development of rigour and quantification in areas of science where formalism had traditionally been looked at with skepticism. The methodologies and interpretations of experimental data that were developed by him and his many collaborators continue to be applied to a wide range of problems of biological and medical significance, especially in the field of protein-protein interactions. A multitude of structure-function studies, recently ranging from the molecular basis of murine olfaction (Portman et al., 2014) through the heterodimeric character of a plant immune receptor (Williams et al., 2014) to a lethal mutation in the laminin alpha-1 gene (Patel et al., 2015), has demonstrated

the importance of the thermodynamic approach to significant molecular biological problems. It can only be anticipated that the influence of Don's work and its application will still be felt many decades into the future.

## Acknowledgment

I express my heartfelt gratitude to Don for 37 prime years of rich scientific comradeship.

## References

- Adams ET Jr, Fujita H (1963) Sedimentation equilibrium in reacting systems. In: Williams JW (ed) Ultracentrifugal analysis in theory and experiment, Academic Press, New York
- Bird RB, Spotz, EL, Hirschfelder JO (1950) The third virial coefficient for non-polar gases. J Chem Phys 18:1395-1402
- Brinkman HC, Hermans JJ (1949) The effect of non-homogeneity of molecular weight on the scattering of light by high polymer solutions. J Chem Phys 17:574-576
- Hill TL (1954) Virial expansions of the osmotic pressure in the Donnan membrane equilibrium, J Chem Phys 22:1251-1252
- Hill TL (1955a) Theory of protein solutions. I. J Chem Phys 23:623-636
- Hill TL (1955b) Theory of protein solutions. II. J Chem Phys 23:2270-2274
- Hill TL (1956a) Osmotic pressure, protein solutions and active transport I. J Am Chem Soc 78:4281-4284
- Hill TL (1956b) On the theory of the Donnan membrane equilibrium. Discuss Faraday Soc 21:31-45
- Hill TL (1957) Theory of solutions. I. J Am Chem Soc 79:4885-4890
- Hill TL (1958) Osmotic pressure, protein solutions and active transport II. J Am Chem Soc 80:2923-2926
- Hill TL (1959) Theory of solutions. II. Osmotic pressure virial expansion and light scattering in two component solutions, J Chem Phys 30:93-97
- Hill TL (1968) Thermodynamics for chemists and biologists, Addison Wesley, Reading, MA
- Hill TL, Chen YD (1973) Theory of aggregation in solution. I. General equations and application to the stacking of bases, nucleosides, etc. Biopolymers 12:1285-1312
- Jacobsen MP, Wills PR, Winzor DJ (1996) Thermodynamic analysis of the effects of small inert cosolutes in the ultracentrifugation of noninteracting proteins. Biochemistry 35:13173-13179

- Kirkwood JG, Buff FP (1951) The statistical mechanical theory of solutions. I. J Chem Phys 19:774-777
- McMillan WG, Mayer JE (1945) The statistical thermodynamics of multicomponent systems. J Chem Phys 13:276-305
- Nichol LW, Ogston AG, Wills PR (1981) Effect of Inert Polymers on Protein Self-Association. FEBS Lett 126:18-20
- Nichol LW, Wills PR, Winzor DJ (1979) An Examination of the Forms of Scatchard Plots of Binding Results Outside Domains of Sigmoidality. J Theor Biol 80:39-50
- Ogston AG, Winzor DJ (1975) Treatment of thermodynamic nonideality in equilibrium studies on associating solutes. J Phys Chem 79:2496-2500.
- Patel TR, Nikodemus D, Besong TMD, et al. (2015) Biophysical analysis of a lethal laminin alpha-1 mutation reveals altered self-interaction. Matrix Biol 49:93-105
- Portman KL, Long J, Carr S et al. (2014) Enthalpy/entropy compensation effects from cavity desolvation underpin broad ligand binding selectivity for rat odorant binding protein 3. Biochemistry. 53:2371-2379
- Scatchard (1946) Physical chemistry of protein solutions. I. Derivation of the equations for the osmotic pressure. J Am Chem Soc 68:2315-2319
- Scott DJ, Wills PR, Winzor DJ (2010) Allowance for the effect of protein charge in the characterization of nonideal solute self-association by sedimentation equilibrium. Biophys Chem 149:83-91
- Stockmayer WH (1950) Light scattering in multi-component systems. J Chem Phys 18:58-61
- Williams SJ, Sohn KH, Wan L, et al. (2014) Structural basis for assembly and function of a heterodimeric plant immune receptor. Science 344:299-303
- Wills PR (1986) Scrapie, ribosomal proteins and biological information (1986) J Theor Biol 122:157-178
- Wills PR, Comper WD, Winzor DJ (1993) Thermodynamic non-ideality in macromolecular solutions: interpretation of virial co-efficients. Arch Biochem Biophys 300:206-212
- Wills PR, Georgalis Y (1981) Concentration Dependence of the Diffusion Coefficient of a Dimerizing Protein: Bovine Pancreatic Trypsin Inhibitor]. Phys Chem 85:3978-3984
- Wills PR, Georgalis Y, Dijk J, Winzor DJ (1995) Measurement of thermodynamic nonideality arising from volume-exclusion between proteins and polymers Biophys Chem 57:37-46
- Wills PR, Hall DR, Winzor DJ (2000) Interpretation of thermodynamic nonideality in sedimentation equilibrium experiments on proteins. Biophys Chem 84:217-225
- Wills PR, Jacobsen MP Winzor DJ (1996) Direct analysis of solute self-association by sedimentation equilibrium. Biopolymers 38:119-130

- Wills PR, Jacobsen MP Winzor DJ (2000) Analysis of sedimentation equilibrium distributions reflecting nonideal macromolecular associations. *Biophys J* 79:2178-2187 (2000)
- Wills PR, Nichol LW, Siezen RJ (1980) The Indefinite Association of Lysozyme; Consideration of Composition-Dependent Activity Coefficients. *Biophys Chem* 11:71-82
- Wills PR, Scott DJ, Winzor DJ (2007) A simpler analysis for the measurement of second virial coefficients by self-interaction chromatography. *Anal Biochem* 371:21-25
- Wills PR, Scott DJ, Winzor DJ (2012) Allowance for effects of thermodynamic nonideality in sedimentation equilibrium distributions reflecting protein dimerization. *Anal Biochem* 422:28-32
- Wills PR, Winzor DJ (1993) Thermodynamic Analysis of 'Preferential Solvation' in Protein Solutions. *Biopolymers* 33:1627-1629
- Wills PR, Winzor DJ (2001) Studies of solute self-association by sedimentation equilibrium: allowance for effects of thermodynamic non-ideality beyond the consequences of nearest-neighbor interactions. *Biophys Chem* 91, 253-262
- Wills PR, Winzor DJ (2002) Exact theory of sedimentation equilibrium made useful. *Progr Colloid Polymer Sci* 119:113-120
- Wills PR, Winzor DJ (2005) Van der Waals phase transition in protein solutions. *Acta Cryst D* 61:832-836
- Wills PR, Winzor DJ (2009) Direct allowance for the effects of thermodynamic nonideality in the quantitative characterization of protein self-association by osmometry. *Biophys Chem* 145:64-71
- Wills PR, Winzor DJ (2011) Allowance for thermodynamic nonideality in the characterization of protein interactions by spectral techniques. *Biophys Chem* 158:21-25
- Winzor DJ, Deszczynski M, Harding SE, Wills PR (2007) Nonequivalence of Second Virial Coefficients from Sedimentation Equilibrium and Light Scattering Studies of Protein Solutions. *Biophys Chem* 128:46-55
- Winzor DJ, Wills PR (1986) Effect of thermodynamic non-ideality on protein interactions: equivalence of interpretations based on excluded volume and preferential solvation. *Biophys Chem* 25:243-251
- Winzor DJ Wills PR (1995) Thermodynamic nonideality of enzyme solutions supplemented with inert solutes: yeast hexokinase revisited. *Biophys Chem* 57:103-110
- Winzor DJ Wills PR (2003) Allowance for thermodynamic non-ideality in the characterization of protein self-association by frontal exclusion chromatography: hemoglobin revisited. *Biophys Chem* 104:345-359
- Winzor DJ, Wills PR (2006) Molecular crowding effects of linear polymers in protein solutions. *Biophys Chem* 119:186-195
- Winzor DJ, Wills PR (2007) Characterization of weak protein dimerization by direct analysis of sedimentation equilibrium distributions: the INVEQ approach. *Anal Biochem* 368:168-177

Woolley P, Wills PR (1985) Excluded-Volume Effect of Inert Macromolecules on the Melting of Nucleic Acids. *Biophys Chem* 22:89-94

**Figure 1 Caption**

Letter from teacher (Terrell Hill) to student (the author) outlining the derivation of the osmotic pressure equation expressed alternatively in terms of molar and molal quantities.