

# Big Heart Data: Advancing Health Informatics through Data Sharing in Cardiovascular Imaging

Avan Suinesiaputra<sup>1</sup>, Pau Medrano-Gracia<sup>1</sup>, Brett R. Cowan<sup>1</sup>, and Alistair A. Young<sup>1</sup>

<sup>1</sup>Dept. of Anatomy with Radiology, University of Auckland, New Zealand

The burden of heart disease is rapidly worsening due to increasing prevalence of obesity and diabetes. Data sharing and open database resources for heart health informatics are important for advancing our understanding of cardiovascular function, disease progression and therapeutics. Data sharing enables valuable information, often obtained at considerable expense and effort, to be re-used beyond the specific objectives of the original study. Many government funding agencies and journal publishers are requiring data re-use, and are providing mechanisms for data curation and archival. Tools and infrastructure are available to archive anonymous data from a wide range of studies, from descriptive epidemiological data to gigabytes of imaging data. Meta-analyses can be performed to combine raw data from disparate studies to obtain unique comparisons or to enhance statistical power. Open benchmark datasets are invaluable for validating data analysis algorithms and objectively comparing results. This review provides a rationale for increased data sharing and surveys recent progress in the cardiovascular domain. We also highlight the potential of recent large cardiovascular epidemiological studies enabling collaborative efforts to facilitate data sharing, algorithms benchmarking, disease modeling and statistical atlases.

*Index Terms*—Cardiac MRI, cardiac atlas, anatomical models, data sharing.

## I. INTRODUCTION

**C**ARDIOVASCULAR disease (CVD) is the world's leading cause of morbidity and mortality. Healthcare costs from CVD are rising due to the increasing prevalence of obesity, diabetes and metabolic syndrome. The results of the Global Burden of Disease Study show that ischemic heart disease was the leading cause of disability-adjusted life years in 2010 [1]. In many countries, obesity is likely to overtake tobacco as the leading risk to health by 2016 [2].

The ability to integrate data from multiple sources and across many variables has significant potential for the evaluation and treatment of patients. These methods enable deeper characterization of a particular patient, and more precise mapping to similar patients in pertinent subpopulations [3]. Efficient characterization of subpopulations requires data harmonization, new analytical algorithms and analysis of heterogeneous data. This approach relies on data sharing infrastructure and high-throughput database resources in large epidemiological studies as well as in small experimental studies, which would otherwise require a large cost or effort to reproduce.

Data sharing for scientific advancement is well established in many fields. In genomics, databases have existed for many years for archiving and curating data [4]. In neuroscience, large repositories are available for specific disease groups or general atlases [5]. Efforts are now being made to facilitate data sharing in the cardiovascular domain [6]. This review highlights the rationale and the need for data sharing in the cardiovascular research community, and outlines current efforts to provide

infrastructure and tools for data sharing. Examples are given by atlas-based analysis of cardiac remodeling and ongoing community efforts to establish benchmarking platforms for cardiac image analysis.

## II. LARGE-SCALE CARDIOVASCULAR POPULATION STUDIES

A large part of our current understanding of multivariable risk factors in the etiology of CVD arose from the Framingham Heart Study, which began in 1948 and is still continuing with over 1,200 publications. The Framingham scores assess cardiovascular risk factors of hypertension, smoking, lipid profile, obesity, diabetes and inactivity; the pathophysiological progression from hypertension to heart failure; the relationship between atrial fibrillation and stroke; and the value of population-based longitudinal studies [7]. With the advent of large-scale databases and data mining methods, it has recently become possible to identify relationships across many different types of information [8].

A wealth of data is now available from non-invasive cardiac imaging examinations. Prospective longitudinal studies derived from large-scale imaging data have been established to investigate the pathogenesis of cardiac diseases [11], [16]. Longitudinal studies, which follow patients over the years, enable scientists to understand the evolution of cardiac disease from sub-clinical manifestations to clinical symptoms. The combination of imaging data with other diagnostic information and biomarkers gives a new rich field of big heart data (defined for the purposes of this review as imaging studies combined with clinical assessments in large study cohorts on an unprecedented scale). These present significant opportunities for new discoveries to reduce the burden of CVD.

Table I lists existing large-scale cardiovascular studies with collection of gigabytes of imaging data. The Multi-Ethnic

The Cardiac Atlas project was supported by Award Number R01HL087773 and R01HL121754 from the National Heart, Lung, and Blood Institute. A. Suinesiaputra, P. Medrano-Gracia, B. R. Cowan and A. A. Young are with the Department of Anatomy with Radiology, Faculty of Medical and Health Sciences, 95 Park Road, University of Auckland, Auckland 1142, New Zealand (e-mails: [a.suinesiaputra; p.medrano; b.cowan; a.young]@auckland.ac.nz).

TABLE I  
MAJOR CARDIOVASCULAR EPIDEMIOLOGICAL STUDIES WITH THE  
INCLUSION OF IMAGING DATA.

Study	Country	Images	Start	Size	Group	Age
MESA [9]	USA	ECG, CT, MRI	2000	6,814	Hispanic, Chinese, White, African- American	45–84
JHS [10]	USA	ECG, CT, MRI	2000	5,302	African- American	21–84
UK Biobank [11]	UK	MRI, ECG, DEXA	2006	100,000	Multi	40–69
CPTP [12]	Canada	MRI	2009	10,000	Multi	35–69
Iceland MI [13]	Iceland	MRI	2004	936	European	67–93
Framingham Offspring [14]	USA	MRI	2002	1,707	Multi	NA
EuroCMR [15]	Europe	MRI	2012	27,781	Multi	47–70

Study of Atherosclerosis (MESA) study was initiated in 2000 to focus on the manifestation of subclinical to clinical CVD before signs and symptoms develop in the heterogeneous population of the US [9]. The study has sampled 6,814 men and women aged 45–84 years old across six centers. The analysis of the 10 year follow-up has recently been completed for around 3,000 participants [17]. The MESA study was the first major population-based study to use cardiac MRI [18]. Data sharing is provided by the MESA study through their ancillary studies and publication protocols.

The UK Biobank [11] is an extensive collection of cardiac data, including questionnaires, physical examinations and biological samples, from 500,000 men and women aged 40–69 in 22 centers across the UK. An imaging enhancement sub-study began in April 2014 with the aim of imaging 6,000 participants during the pilot phase. If successful, 100,000 participants will be imaged with MRI over a 5–6 year period, which will be the largest prospective cardiac imaging dataset worldwide. Imaging modalities include cardiac MRI, abdominal MRI, brain MRI, carotid ultrasound, and DEXA.

The Jackson Heart Study (JHS) was designed to determine the root causes of CVD in 5,302 African-American individuals aged 21–84 years living in the southeastern USA (Jackson, Mississippi) [10], [19], [20]. This group experiences increased mortality from CVD as well as higher incidence of hypertension, obesity and diabetes. A unique aspect of JHS was the association of neighborhood disadvantages in terms of economic, sociocultural, behavioral, dietary and physical activity measures with cardiometabolic risk factors [21], [22]. The JHS provides a rare insight into the interactions between genotype and phenotype in a high-risk population. In terms of imaging data, the JHS has collected 3,000 CT and MRI examinations containing heart function and calcium scores [23].

Other studies such as the Canadian Partnership for Tomorrow Project (CPTP) with 10,000 participants, ICELAND MI with 936 patients with myocardial infarction, and the

Framingham Offspring study with 1,707 follow-up cardiac MRI scans added more imaging data for cardiac research. While not a study per se, the EuroCMR registry has the main goal to evaluate the prognostic potential of cardiac MRI as well as its cost-effectiveness. More than 27,000 consecutive patients have been enrolled from 57 centers in 15 countries. Similarly, the Global CMR registry has been established to collate MRI patient data from around the world with 44,000 cases contributed to date.

### III. SHARING MEDICAL DATA

With the ability to mine multidimensional relationships in diverse datasets, sharing patient medical records across clinical providers and researchers becomes a key factor to improve quality of care, to reduce healthcare costs and to minimize human errors [24]. However, sharing medical data remains a big challenge due to the extremely granular nature of healthcare data. Hospitals, imaging centers, healthcare institutions and clinics must enforce strict guidelines to uphold patient data privacy and security. Therefore, central to enable medical data sharing is a secured protection of patient data, such as mandated by the HIPAA (Health Insurance Portability and Accountability Act) in the US or the Data Protection Act in the UK. Ensuring patient data privacy is vital to sharing medical records to a wider community.

In general, there are three entities in data sharing: *data contributors*, *organizers* and *users* (see Fig. 1). The data contributor is a provider who wants to release their data to a wider research community without breaching agreements with patients and other clinical interests. Data contributors must ensure informed consent is obtained, which is compatible with data sharing, in conjunction with local ethics committee or institutional review board (IRB) approval. They can only contribute anonymized or de-identified data. The de-identification process removes any protected health information that can be used to identify an individual, e.g. name, dates (except




Fig. 1. Three components of medical data sharing: data contributor, organizer and user. The data contributor controls the acceptance of user request, while the organizer manages, stores and also enriches the original clinical data with derived analyses. This figure was adapted from [6].

TABLE II  
EXISTING MEDICAL DATA SHARING INFRASTRUCTURES THAT INCLUDE CARDIAC DATA

Name	Domain	Accessibility	Studies (size)	Data type	URL
CAP	Cardiac	Registration Guest	Asymptomatic (2,450) Myocardial infarction (465) Normal human heart atlas (1) Mixed normal, hypertrophy, heart failure (45)	MRI Finite element Contours Binary images	www.cardiacatlas.org
AMDB	Cardiac	Registration	Unspecified	Geometrical models	amdb.isd.kcl.ac.uk
iDASH	Breast Sequencing data Cardiac Colon Lung Physical motion Lab tests General medicine	Registration Guest	Breast MRI (153) Clinical logs (62) Radiology reports (2,363) CT colonography (103) Diabetes mellitus data (17) Laboratory test data (13,000) Lung CT (398+) Coronary artery disease data (20,000) ICU record (53) Motion sensor data (16)	R data MRI Text files CT GPS data CSV files JPEG images	idash.ucsd.edu
CVRG	Cardiac	Open	Normal canine heart (7) Failing canine heart (5) Normal human (1) Canine heart atlas (1) Ischemic human CT (13) Non-ischemic human CT (12)	MRI CT	cvrgrid.org
PhysioNet	Cardiac Brain Biomedical signals	Open	European ST-T database (79) Long-term ST database (43) Arrhythmia database (23) Noise stress test (14) Heart rate database (10) Congestive heart failure (15) Partial epilepsy (7) QT database (100)	ECG	www.physionet.org
VIP	Various organs	Registration	Unspecified	Simulated data	www.creatis.insa-lyon.fr/vip

for year), social security, location or other unique identifiers. These can be stored in the metadata, concealed as filename, or even hidden in clinical narratives [25]. Several automated de-identification algorithms are available (see a survey in [26]).

To ensure that IRB and study requirements are met, data contributors are usually involved in a user review process to decide whether to grant or to deny a request for data access. Alternatively, data contributors can determine specific types of research activities. For example, the left ventricular consensus segmentation project [27] uses a subset from the DETERMINE cohort [28], where the data contributor has given approval to use 200 MRI cases of patients with myocardial infarction specifically for the development of automatic segmentation algorithms. In another framework called PCARE [29], patients also act as data contributor, who control how their data are being shared to unaffiliated healthcare organizations.

For the organizer who wants to set up data sharing infrastructure, the main challenges are to ensure the integrity of the data and to establish seamless integration, storage and management of a massive amount of heterogeneous data, which may include images, laboratory tests, diagnostic records, biological and physiological models. The information technology to share multi-dimensional data over the internet has matured in the last decade. Several data sharing infrastructure alternatives have been proposed in various domains, such as [6], [29]–[31].

The organizer may also be involved in a process to enrich the original patient data to make the data more useful for broader research activities. Such data enrichment activities

include ontological annotations, manual labeling, model fitting and population analysis (see Fig. 1).

User responsibility can be defined by a data distribution agreement, which dictates the specific type of research, commitment to secure the data within their host computer, and a potential intellectual property agreement. All conditions or restrictions on data use should be specified in the data distribution agreement.

#### IV. EXISTING CARDIAC DATA SHARING

In the cardiac domain, several efforts have been initiated to establish infrastructures for data sharing by providing anonymized baseline examinations, imaging data and other derived computations, such as anatomical models and statistical shape analysis. Table II summarizes existing medical data sharing infrastructures, which also include cardiovascular data.

Established in 2010, the Cardiac Atlas Project (CAP) is a worldwide consortium to host large cardiac image data with derived finite element models of the heart and associated diagnostic information [6]. Over 3,000 cardiac MRI cases have been contributed to the database, which are being used by more than 20 research groups worldwide for various research activities. CAP is a registered ancillary study of MESA and has developed methods to translate results between MESA and other studies using atlas-based bias correction methods [32]. Patient-specific models of the heart provide a standard coordinate system, which map the heart according to anatomical location. CAP has developed methods to pool data from different sources in a standardized manner, and

to correct bias arising from imaging or analysis protocol. CAP is endorsed by the Society of Cardiovascular Magnetic Resonance, which maintains an upload site where cases can be contributed to the atlas project [33].

The Anatomical Models Database (AMDB) is a web accessible framework to share and reuse cardiovascular models [30], [31]. AMDB stores various cardiac geometry models, accessible for any researchers to perform a simulation or a benchmarking study in cardiac electrophysiology and mechanics. The aim of AMDB (formerly known as euHeart) is to allow multi-scale computational modeling of the heart [34]. AMDB employs a web service tool that personalizes a geometric heart model given binary images of the heart and cardiac parameters [35].

The iDASH (integrated data analysis, anonymization and sharing) is a general framework to provide a scalable tool to share and access medical data, including cardiac diseases [36]. The iDASH stores and shares heterogeneous data from multiple domains (see Table II). A specific infrastructure that shares only ECG signals is provided by PhysioNet. Currently, PhysioNet stores and freely shares large scale de-identified recorded physiologic signals, time series and other related biomedical data [37]. These are healthy subjects and patients with various diseases, including sudden cardiac death, congestive heart failures, sleep apnea, aging, epilepsy and gait disorder. The CardioVascular Research Grid (CVRG) provides an infrastructure to securely and seamlessly access complex data of cardiovascular studies, including an automated ontological labeling of anatomical differences [38]. CVRG is currently establishing an easy access platform with cloud-based and browser-based tools without the need to install complex software. The Virtual Imaging Platform (VIP) is another openly accessible online data sharing platform, which focuses more on computationally extensive biomedical simulation processes [39].

## V. APPLICATION: POPULATION-BASED CARDIAC REMODELING

The human heart is continuously remodeling (changing shape) in response to pathology, aging, environmental and genetic factors. Cardiac remodeling can be maladaptive when linked to heart failure progression [40], [41], but remodeling can be adaptive during normal growth or intensive physical exercises. Even in the early stages of heart failure, adaptive remodeling can be observed because the heart maintains its function in spite of pressure or volume overloading in the acute phase of cardiac injury [42]. There is a transition from adaptive to maladaptive remodeling in the progression of heart disease, but when and how this transition occurs still remains unknown [43]. As illustrated in Fig. 2, population-based studies that combine cardiac shape, function and other clinical data can characterize these remodeling processes.

As the heart remodels, its geometry, mass, composition and volume changes. The shape of the heart can become less elliptical and more spherical [44]. Ventricular sphericity (width to height ratio) has been observed in symptomatic patients and associated with decreased survival [45] and adverse remodeling [46]. Increased LV chamber dimension [47], lower systolic

dimension change [48] and hypertrophy [49] have also been observed in asymptomatic individuals. Fig. 3 shows how LV size and sphericity were increased from normal volunteers to patients with heart failure in a subset of cases from CAP.

The understanding of cardiac remodeling processes is particularly important in quantifying effects of treatment and reverse remodeling [50]. Several studies have been actively investigating how remodeling occurs in the population by using statistical shape and function tools [51]–[56]. For instance, principal component analysis (PCA) on the cardiac shapes of 102 adults born preterm (30 weeks) and 102 age matched healthy volunteers have shown different shape indices in adults who were born preterm, compared with normal term controls [53]. A statistical cardiac atlas built from 1991 asymptomatic individuals has shown a relationship between traditional risk factors (hypertension, smoking, sex and diabetes) with LV remodeling [56]. These studies have demonstrated the power of geometrical heart shape models to identify cardiac remodeling in certain sub-populations or in later life.

## VI. APPLICATION: BENCHMARKING

A major advantage of data sharing is the availability of standardized datasets that can be used to develop, validate and compare (benchmark) new automated algorithms. Benchmarking is important to evaluate the relative efficacy of the growing number of proposed automated methods. Objective comparisons are more difficult to perform if each published method presents results using private data and incompatible platforms.

Community efforts have been made to provide platforms for benchmarking on existing and emerging algorithms through a series of “challenges”. A left ventricular segmentation challenge was held in 2009 using 45 cases from a mixed patient dataset (normal, heart failure, myocardial infarction and hypertrophy) [57]. These data are in the public domain and can be accessed directly via the Cardiac Atlas Project website. A similar challenge was run in 2011 with more cases (200 patients with myocardial infarction) [58], which subsequently established a platform for collaboration between research groups to define a common consensus ground truth for the myocardial segmentation [27]. Other benchmarking frameworks utilizing cardiac data sharing include cardiac motion [59], segmentation of scar tissue in the left atrium [60], and automated extraction of centerlines in coronary arteries [61]. Future challenges and benchmarking studies are planned, including cardiac mechanics, electrophysiology and correcting breathing motion artifacts in perfusion MR images.

## VII. CHALLENGES

The main challenge in big heart data studies lies primarily on the willingness of providers to release data anonymously to a wider community. Fortunately, there is an ongoing progression towards open access data required by national funding agencies for new research projects, such as the National Institutes of Health in the US and the National Health Services in the UK. Reproducibility of research outcomes is




Fig. 2. Cardiac remodeling process over time (green = adaptive remodeling due to growth, red = maladaptive remodeling). Individual dots illustrate image acquisition. The availability of big heart data may enable prediction of remodeling and risk stratification of individuals at risk.

the main driving force of this research direction. The onus is now on organizers to provide robust and user-friendly data sharing platforms. They must guarantee to both providers and the general public that patient data have been de-identified properly, data integrity are maintained, and data dissemination occurs in the most secure way.

Standardization is a key technical challenge in data sharing. The DICOM standard for storing pixel and non-pixel data (patient information, 3D image geometry, examination records, acquisition parameters, etc.) can change from vendor to vendor or even software version. Picture Archiving and Communication System (PACS) allows radiographic images being stored and transferred electronically within and between hospitals. Open source PACS architectures, such as dcm4chee [62], enable quick deployment of a web-accessible imaging data server for public use. Data standardization is even more difficult for non-radiographic image data. Initial efforts have been proposed, including the standardization of a modeling language (FieldML) to link different organs and tissues [63], an open electrocardiogram (ECG) format [64] and cardiac electrophysiology (EP) key data elements [65].

## VIII. FUTURE PERSPECTIVES

### A. Novel geometry-based cardiovascular risk predictors

Of all various pathways to end-stage CVD, the primary preventive measure is to reduce risk factors and to prevent early disease progression. Improving major risk factors has led to considerable reduction in morbidity and mortality due to CVD, but yet CVD remains the leading cause of death and




Fig. 3. Multivariate map constructed from the first two principal components of left-ventricular shape at end-systole (ES). Heart failure (HF) cases with and without infarct show elevated size and sphericity with these measures. However, hypertrophic cases are similar in these measures to normal volunteers. Data were taken from public domain Sunnybrook Cardiac Dataset available from the Cardiac Atlas Project.

disability in developed countries. Traditional CVD risk assessment currently disregards the geometrical shape of the heart, which leads to incomplete information about how the heart remodels over time in a matched sub-clinical group, global and regional functional parameters and other biomechanical properties. Early diagnosis and risk stratification of individuals at risk can be improved by augmenting current risk factors predictors [66], a process which can be accelerated by sharing cardiac data.

### B. Disease modeling

The availability of cardiac models enables development of computer-aided diagnosis applications for cardiac diseases [32], [67]. Subtle differences in cardiac motion, which are inherently difficult to assess, can be objectively quantified by a statistical model of cardiac motion in the population. Such applications will allow better tracking and monitoring of the progression of a cardiac disease over time. This becomes essential in a rapidly progressive disease such as pulmonary arterial hypertension, where regular assessment is needed before tailoring the treatment in each patient [68].

### C. Personalization

Collecting individual data gains statistical knowledge about the population; a valuable resource to be brought back to individuals [69] by means of a personalized treatment [34], [70], [71]. Preliminary studies in this frontier include the prediction of fiber orientation in the left ventricle [72], heart

geometry personalization [73], [74], predicting cardiac resynchronization response [75] and generating an artificial patient-specific model for pre-surgical planning [76]. Personalized medicine will gain more attention in the future and will be more tightly connected with the availability of big heart data.

## IX. CONCLUSION

Data sharing is becoming essential to the further development of the field of cardiovascular research. Large databases are now being developed which bring together disparate types of data and enable multi-variate greater-depth analysis of information on each patient. In particular, population-based tools have proven useful for comparing one individual's anatomy to a population or between sub-populations, and in the detection, quantification and monitoring the disease progression. Data-sharing initiatives will aid in the validation and further automation of data analysis methods.

## REFERENCES

- [1] C. J. L. Murray, T. Vos, R. Lozano *et al.*, "Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010," *Lancet*, vol. 380, no. 9859, pp. 2197-223, Dec 2012.
- [2] "Health Loss in New Zealand: A report from the New Zealand Burden of Diseases, Injuries and Risk Factors Study, 2006-2016," Ministry of Health, Wellington, 2013.
- [3] I. S. Kohane, "The twin questions of personalized medicine: who are you and whom do you most resemble?" *Genome Med*, vol. 1, no. 1, p. 4, 2009.
- [4] T. P. Sneddon, P. Li, and S. C. Edmunds, "GigaDB: announcing the GigaScience database," *Gigascience*, vol. 1, no. 1, p. 11, 2012.
- [5] S. Mori, K. Oishi, A. V. Faria, and M. I. Miller, "Atlas-based neuroinformatics via MRI: harnessing information from past clinical cases and quantitative image analysis for patient care," *Annu Rev Biomed Eng*, vol. 15, pp. 71-92, 2013.
- [6] C. G. Fonseca, M. Backhaus, D. A. Bluemke *et al.*, "The Cardiac Atlas Project—an imaging database for computational modeling and statistical atlases of the heart," *Bioinformatics*, vol. 27, no. 16, pp. 2288-95, Aug 2011.
- [7] W. B. Kannel, T. R. Dawber, A. Kagan, N. Revotskie, and J. Stokes III, "Factors of risk in the development of coronary heart disease—six year follow-up experience. The Framingham Study," *Ann Intern Med*, vol. 55, pp. 33-50, Jul 1961.
- [8] T. B. Murdoch and A. S. Detsky, "The inevitable application of big data to health care," *JAMA*, vol. 309, no. 13, pp. 1351-2, Apr 2013.
- [9] D. E. Bild, D. A. Bluemke, G. L. Burke *et al.*, "Multi-ethnic study of atherosclerosis: objectives and design," *Am J Epidemiol*, vol. 156, no. 9, pp. 871-81, Nov 2002.
- [10] H. A. Taylor, Jr, J. G. Wilson, D. W. Jones *et al.*, "Toward resolution of cardiovascular health disparities in African Americans: design and methods of the Jackson Heart Study," *Ethn Dis*, vol. 15, no. 4 Suppl 6, pp. S6-4-17, 2005.
- [11] S. E. Petersen, P. M. Matthews, F. Bamberg *et al.*, "Imaging in population science: cardiovascular magnetic resonance in 100,000 participants of UK Biobank - rationale, challenges and approaches," *J Cardiovasc Magn Reson*, vol. 15, p. 46, 2013.
- [12] M. J. Borugian, P. Robson, I. Fortier *et al.*, "The Canadian Partnership for Tomorrow Project: building a pan-Canadian research platform for disease prevention," *CMAJ*, vol. 182, no. 11, pp. 1197-201, Aug 2010.
- [13] E. B. Schelbert, J. J. Cao, S. Sigurdsson *et al.*, "Prevalence and prognosis of unrecognized myocardial infarction determined by cardiac magnetic resonance in older adults," *JAMA*, vol. 308, no. 9, pp. 890-6, Sep 2012.
- [14] M. L. Chuang, P. Gona, G. L. T. F. Hautvast *et al.*, "CMR reference values for left ventricular volumes, mass, and ejection fraction using computer-aided analysis: the Framingham Heart Study," *J Magn Reson Imaging*, vol. 39, no. 4, pp. 895-900, Apr 2014.
- [15] O. Bruder, A. Wagner, M. Lombardi *et al.*, "European Cardiovascular Magnetic Resonance (EuroCMR) registry—multi national results from 57 centers in 15 countries," *J Cardiovasc Magn Reson*, vol. 15, p. 9, 2013.
- [16] N. D. Wong, "Epidemiological studies of CHD and the evolution of preventive cardiology," *Nat Rev Cardiol*, vol. 11, no. 5, pp. 276-89, May 2014.
- [17] D. E. Bild, R. McClelland, J. D. Kaufman *et al.*, "Ten-year trends in coronary calcification in individuals without clinical cardiovascular disease in the multi-ethnic study of atherosclerosis," *PLoS One*, vol. 9, no. 4, p. e94916, 2014.
- [18] D. A. Bluemke, R. A. Kronmal, J. A. C. Lima *et al.*, "The relationship of left ventricular mass and geometry to incident cardiovascular events: the MESA (Multi-Ethnic Study of Atherosclerosis) study," *J Am Coll Cardiol*, vol. 52, no. 25, pp. 2148-55, Dec 2008.
- [19] J. Harman, E. R. Walker, V. Charbonneau *et al.*, "Treatment of hypertension among African Americans: the Jackson Heart Study," *J Clin Hypertens (Greenwich)*, vol. 15, no. 6, pp. 367-74, Jun 2013.
- [20] H. A. Taylor, Jr, "The Jackson Heart Study: an overview," *Ethn Dis*, vol. 15, no. 4 Suppl 6, pp. S6-1-3, 2005.
- [21] T. J. Payne, S. B. Wyatt, T. H. Mosley *et al.*, "Sociocultural methods in the Jackson Heart Study: conceptual and descriptive overview," *Ethn Dis*, vol. 15, no. 4 Suppl 6, pp. S6-38-48, 2005.
- [22] C. R. Clark, M. J. Ommerborn, D. A. Hickson *et al.*, "Neighborhood disadvantage, neighborhood safety and cardiometabolic risk factors in African Americans: biosocial associations in the Jackson Heart study," *PLoS One*, vol. 8, no. 5, p. e63254, 2013.
- [23] J. J. Carr, "The revolution in risk assessment and disease detection made possible with non-invasive imaging: implications for population science," *Ethn Dis*, vol. 22, no. 3 Suppl 1, pp. S1-24-7, 2012.
- [24] A. E. Flanders, "Medical image and data sharing: are we there yet?" *Radiographics*, vol. 29, no. 5, pp. 1247-51, 2009.
- [25] M. Li, D. Carrell, J. Aberdeen, L. Hirschman, and B. A. Malin, "De-identification of clinical narratives through writing complexity measures," *Int J Med Inform*, vol. 83, no. 10, pp. 750-67, Oct 2014.
- [26] A. Gkoulalas-Divanis, G. Loukides, and J. Sun, "Publishing data from electronic health records while preserving privacy: a survey of algorithms," *J Biomed Inform*, vol. 50, pp. 4-19, Aug 2014.
- [27] A. Suinesiaputra, B. R. Cowan, A. O. Al-Agamy *et al.*, "A collaborative resource to build consensus for automated left ventricular segmentation of cardiac MR images," *Med Image Anal*, vol. 18, no. 1, pp. 50-62, Jan 2014.
- [28] A. H. Kadish, D. Bello, J. P. Finn *et al.*, "Rationale and design for the Defibrillators to Reduce Risk by Magnetic Resonance Imaging Evaluation (DETERMINE) trial," *J Cardiovasc Electrophysiol*, vol. 20, no. 9, pp. 982-7, Sep 2009.
- [29] Y. Ge, D. K. Ahn, B. Unde, H. D. Gage, and J. J. Carr, "Patient-controlled sharing of medical imaging data across unaffiliated healthcare organizations," *J Am Med Inform Assoc*, vol. 20, no. 1, pp. 157-63, Jan 2013.
- [30] D. Gianni, S. McKeever, T. Yu *et al.*, "Sharing and reusing cardiovascular anatomical models over the Web: a step towards the implementation of the virtual physiological human project," *Philos Trans A Math Phys Eng Sci*, vol. 368, no. 1921, pp. 3039-56, Jun 2010.
- [31] E. Kerfoot, P. Lamata, S. Niederer, R. Hose, J. Spaan, and N. Smith, "Share and enjoy: anatomical models database—generating and sharing cardiovascular model data using web services," *Med Biol Eng Comput*, vol. 51, no. 11, pp. 1181-90, Nov 2013.
- [32] P. Medrano-Gracia, B. R. Cowan, D. A. Bluemke *et al.*, "Atlas-based analysis of cardiac shape and function: correction of regional shape bias due to imaging protocol for population studies," *J Cardiovasc Magn Reson*, vol. 15, p. 80, 2013.
- [33] Society of Cardiovascular Magnetic Resonance, "CAP Website." [Online]. Available: <http://www.scmr.org/Research/cap-dicom-transfer.html>
- [34] N. Smith, A. de Vecchi, M. McCormick *et al.*, "euHeart: personalized and integrated cardiac care using patient-specific cardiovascular modelling," *Interface Focus*, vol. 1, no. 3, pp. 349-64, Jun 2011.
- [35] P. Lamata, S. Niederer, D. Nordsletten *et al.*, "An accurate, fast and robust method to generate patient-specific cubic Hermite meshes," *Med Image Anal*, vol. 15, no. 6, pp. 801-13, Dec 2011.
- [36] L. Ohno-Machado, V. Bafna, A. A. Boxwala *et al.*, "iDASH: integrating data for analysis, anonymization, and sharing," *J Am Med Inform Assoc*, vol. 19, no. 2, pp. 196-201, 2012.
- [37] A. L. Goldberger, L. A. Amaral, L. Glass *et al.*, "PhysioBank, PhysioToolkit, and PhysioNet: components of a new research resource for complex physiologic signals," *Circulation*, vol. 101, no. 23, pp. E215-20, Jun 2000.
- [38] S. Steinert-Threlkeld, S. Ardekani, J. L. V. Mejino *et al.*, "Ontological labels for automated location of anatomical shape differences," *J Biomed Inform*, vol. 45, no. 3, pp. 522-7, Jun 2012.



- [39] T. Glatard, C. Lartizien, B. Gibaud *et al.*, "A virtual imaging platform for multi-modality medical image simulation," *IEEE Trans Med Imaging*, vol. 32, no. 1, pp. 110–8, Jan 2013.
- [40] M. A. Konstam, D. G. Kramer, A. R. Patel, M. S. Maron, and J. E. Udelson, "Left ventricular remodeling in heart failure: current concepts in clinical significance and assessment," *JACC Cardiovasc Imaging*, vol. 4, no. 1, pp. 98–108, Jan 2011.
- [41] M. G. Sutton and N. Sharpe, "Left ventricular remodeling after myocardial infarction: pathophysiology and therapy," *Circulation*, vol. 101, no. 25, pp. 2981–8, Jun 2000.
- [42] J. N. Cohn, R. Ferrari, and N. Sharpe, "Cardiac remodeling—concepts and clinical implications: a consensus paper from an international forum on cardiac remodeling. Behalf of an International Forum on Cardiac Remodeling," *J Am Coll Cardiol*, vol. 35, no. 3, pp. 569–82, Mar 2000.
- [43] J. S. Burchfield, M. Xie, and J. A. Hill, "Pathological ventricular remodeling: mechanisms: part 1 of 2," *Circulation*, vol. 128, no. 4, pp. 388–400, Jul 2013.
- [44] G. F. Mitchell, G. A. Lamas, D. E. Vaughan, and M. A. Pfeffer, "Left ventricular remodeling in the year after first anterior myocardial infarction: a quantitative analysis of contractile segment lengths and ventricular shape," *J Am Coll Cardiol*, vol. 19, no. 6, pp. 1136–44, May 1992.
- [45] S. P. Wong, J. K. French, A.-M. Lydon, S. O. M. Manda, W. Gao, N. G. Ashton, and H. D. White, "Relation of left ventricular sphericity to 10-year survival after acute myocardial infarction," *Am J Cardiol*, vol. 94, no. 10, pp. 1270–5, Nov 2004.
- [46] H. F. J. Mannaerts, J. A. van der Heide, O. Kamp, M. G. Stoel, J. Twisk, and C. A. Visser, "Early identification of left ventricular remodelling after myocardial infarction, assessed by transthoracic 3D echocardiography," *Eur Heart J*, vol. 25, no. 8, pp. 680–7, Apr 2004.
- [47] R. S. Vasan, M. G. Larson, E. J. Benjamin, J. C. Evans, and D. Levy, "Left ventricular dilatation and the risk of congestive heart failure in people without myocardial infarction," *N Engl J Med*, vol. 336, no. 19, pp. 1350–5, May 1997.
- [48] M. S. Lauer, J. C. Evans, and D. Levy, "Prognostic implications of subclinical left ventricular dilatation and systolic dysfunction in men free of overt cardiovascular disease (the Framingham Heart Study)," *Am J Cardiol*, vol. 70, no. 13, pp. 1180–4, Nov 1992.
- [49] W. B. Kannel, D. Levy, and L. A. Cupples, "Left ventricular hypertrophy and risk of cardiac failure: insights from the Framingham Study," *J Cardiovasc Pharmacol*, vol. 10 Suppl 6, pp. S135–40, 1987.
- [50] M. A. Konstam, J. E. Udelson, and N. Sharpe, "Prevention and reversal of left ventricular remodeling: summation," *J Card Fail*, vol. 8, no. 6 Suppl, pp. S506–11, Dec 2002.
- [51] N. Duchateau, M. De Craene, G. Piella *et al.*, "A spatiotemporal statistical atlas of motion for the quantification of abnormal myocardial tissue velocities," *Med Image Anal*, vol. 15, no. 3, pp. 316–28, Jun 2011.
- [52] C. Hoogendoorn, N. Duchateau, D. Sánchez-Quintana *et al.*, "A high-resolution atlas and statistical model of the human heart from multislice CT," *IEEE Trans Med Imaging*, vol. 32, no. 1, pp. 28–44, Jan 2013.
- [53] A. J. Lewandowski, D. Augustine, P. Lamata *et al.*, "Preterm heart in adult life: cardiovascular magnetic resonance reveals distinct differences in left ventricular mass, geometry, and function," *Circulation*, vol. 127, no. 2, pp. 197–206, Jan 2013.
- [54] H. Lombaert, J.-M. Peyrat, P. Croisille *et al.*, "Human atlas of the cardiac fiber architecture: study on a healthy population," *IEEE Trans Med Imaging*, vol. 31, no. 7, pp. 1436–47, Jul 2012.
- [55] K. McLeod, C. Seiler, M. Sermesant, and X. Pennec, "Spatio-temporal dimension reduction of cardiac motion for group-wise analysis and statistical testing," *Med Image Comput Comput Assist Interv*, vol. 16, no. Pt 2, pp. 501–8, 2013.
- [56] P. Medrano-Gracia, B. R. Cowan, B. Ambale-Venkatesh *et al.*, "Left ventricular shape variation in asymptomatic populations: the multi-ethnic study of atherosclerosis," *J Cardiovasc Magn Reson*, vol. 16, no. 1, p. 56, Jul 2014.
- [57] Sunnybrook Health Sciences Centre, "Cardiac MR Left Ventricle Segmentation Challenge." [Online]. Available: [http://smial.sri.utoronto.ca/LV\\_Challenge/Home.html](http://smial.sri.utoronto.ca/LV_Challenge/Home.html)
- [58] A. Suinesiaputra, B. R. Cowan, J. P. Finn *et al.*, "Left Ventricular Segmentation Challenge from Cardiac MRI: A Collation Study," in *Statistical Atlases and Computational Models of the Heart. Imaging and Modelling Challenges*, ser. LNCS, vol. 7085. Springer, 2012, pp. 88–97.
- [59] C. Tobon-Gomez, M. De Craene, K. McLeod *et al.*, "Benchmarking framework for myocardial tracking and deformation algorithms: an open access database," *Med Image Anal*, vol. 17, no. 6, pp. 632–48, Aug 2013.
- [60] R. Karim, R. J. Housden, M. Balasubramaniam *et al.*, "Evaluation of current algorithms for segmentation of scar tissue from late gadolinium enhancement cardiovascular magnetic resonance of the left atrium: an open-access grand challenge," *J Cardiovasc Magn Reson*, vol. 15, p. 105, 2013.
- [61] M. Schaap, C. T. Metz, T. van Walsum *et al.*, "Standardized evaluation methodology and reference database for evaluating coronary artery centerline extraction algorithms," *Med Image Anal*, vol. 13, no. 5, pp. 701–14, Oct 2009.
- [62] P. Welter, C. Hocken, T. M. Deserno, C. Grouls, and R. W. Günther, "Workflow management of content-based image retrieval for CAD support in PACS environments based on IHE," *Int J Comput Assist Radiol Surg*, vol. 5, no. 4, pp. 393–400, Jul 2010.
- [63] R. D. Britten, G. R. Christie, C. Little *et al.*, "FieldML, a proposed open standard for the Physiome project for mathematical model representation," *Med Biol Eng Comput*, vol. 51, no. 11, pp. 1191–207, Nov 2013.
- [64] R. R. Bond, D. D. Finlay, C. D. Nugent, and G. Moore, "A review of ECG storage formats," *Int J Med Inform*, vol. 80, no. 10, pp. 681–97, Oct 2011.
- [65] American College of Cardiology/American Heart Association Task Force on Clinical Data Standards (ACC/AHA/HRS Writing Committee to Develop Data Standards on Electrophysiology), A. E. Buxton, H. Calkins *et al.*, "ACC/AHA/HRS 2006 key data elements and definitions for electrophysiological studies and procedures: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Data Standards (ACC/AHA/HRS Writing Committee to Develop Data Standards on Electrophysiology)," *Circulation*, vol. 114, no. 23, pp. 2534–70, Dec 2006.
- [66] R. Hinojar, R. Botnar, J. C. Kaski *et al.*, "Individualized cardiovascular risk assessment by cardiovascular magnetic resonance," *Future Cardiol*, vol. 10, no. 2, pp. 273–89, Mar 2014.
- [67] A. Suinesiaputra, A. F. Frangi, T. A. M. Kaandorp *et al.*, "Automated detection of regional wall motion abnormalities based on a statistical model applied to multislice short-axis cardiac MR images," *IEEE Trans Med Imaging*, vol. 28, no. 4, pp. 595–607, Apr 2009.
- [68] J.-L. Vachiéry, P. Yerly, and S. Huez, "How to detect disease progression in pulmonary arterial hypertension," *Eur Respir Rev*, vol. 21, no. 123, pp. 40–7, Mar 2012.
- [69] A. A. Young and A. F. Frangi, "Computational cardiac atlases: from patient to population and back," *Exp Physiol*, vol. 94, no. 5, pp. 578–96, May 2009.
- [70] C. Stratz, M. Amann, D. D. Berg *et al.*, "Novel biomarkers in cardiovascular disease: research tools or ready for personalized medicine?" *Cardiol Rev*, vol. 20, no. 3, pp. 111–7, 2012.
- [71] M. Ouzounian, D. S. Lee, A. O. Gramolini *et al.*, "Predict, prevent and personalize: Genomic and proteomic approaches to cardiovascular medicine," *Can J Cardiol*, vol. 23 Suppl A, pp. 28A–33A, Aug 2007.
- [72] K. Lekadir, C. Hoogendoorn, M. Pereanez *et al.*, "Statistical personalization of ventricular fiber orientation using shape predictors," *IEEE Trans Med Imaging*, vol. 33, no. 4, pp. 882–90, Apr 2014.
- [73] P. Lamata, M. Sinclair, E. Kerfoot *et al.*, "An automatic service for the personalization of ventricular cardiac meshes," *J R Soc Interface*, vol. 11, no. 91, p. 20131023, Feb 2014.
- [74] M. W. Krueger, W. H. W. Schulze, K. S. Rhode *et al.*, "Towards personalized clinical in-silico modeling of atrial anatomy and electrophysiology," *Med Biol Eng Comput*, vol. 51, no. 11, pp. 1251–60, Nov 2013.
- [75] M. Sermesant, R. Chabiniok, P. Chinchapatnam *et al.*, "Patient-specific electromechanical models of the heart for the prediction of pacing acute effects in CRT: a preliminary clinical validation," *Med Image Anal*, vol. 16, no. 1, pp. 201–15, Jan 2012.
- [76] J. Weese, A. Groth, H. Nickisch, H. Barschdorf *et al.*, "Generating anatomical models of the heart and the aorta from medical images for personalized physiological simulations," *Med Biol Eng Comput*, vol. 51, no. 11, pp. 1209–19, Nov 2013.