







**CDMTCS
Research
Report
Series**




**Topologies for Finite Words:
Compatibility with the CANTOR
Topology**



Ludwig Staiger
Martin-Luther-Universität
Halle-Wittenberg



CDMTCS-568
November 2022



Centre for Discrete Mathematics and
Theoretical Computer Science

Topologies for Finite Words: Compatibility with the CANTOR Topology

Ludwig Staiger

Institut für Informatik,
Martin-Luther-Universität Halle-Wittenberg
D-06099 Halle, Germany
`staiger@informatik.uni-halle.de`

Abstract

Infinite words are often considered as limits of finite words. As topological methods have been proved to be useful in the theory of ω -languages it seems to be providing to include finite and infinite words into one (topological) space. In most cases **this results** in a poor topological structure induced on the subspace of finite words.

In the present paper we investigate the possibility to link topologies in the space of finite words with a topology in the space of infinite words via a natural mapping. A requirement in this linking of topologies consists in the compatibility of the topological properties (openness, closedness etc) of images with preimages and vice versa.

Here we show that choosing for infinite words the natural topology of the CANTOR space and the δ -limit as linking mapping there are several natural topologies on the space of finite words compatible with the topology of the CANTOR space. It is interesting to observe that besides the well-known prefix topology there are at least two more whose origin is from language theory—centers and super-centers of languages.

We show that several of these topologies on the space of finite words fit into a class of \mathcal{L} -topologies and exhibit their special properties w.r.t. to the compatibility with the CANTOR topology.

Contents

1	Introduction	2
2	Preliminaries	4
2.1	Notation	4
2.2	General topology	5
3	Topologies for Words and ω-words	6
3.1	The CANTOR topology on X^ω	7
3.2	The prefix topology \mathcal{T}_p on X^*	7
3.3	Topologies on $X^* \cup X^\omega$	8
4	Linking Topologies on X^* to CANTOR Topology	9
4.1	Compatibility of topologies	9
4.2	(σ, δ) -subsets of X^*	10
5	\mathcal{L}-topologies	11
5.1	The operator Anf	12
5.2	\mathcal{L} -topologies	13
6	Strongly Compatible Topologies on X^*	14
6.1	Center and Supercenter topologies	14
6.2	Strong compatibility - characterisation	16
6.3	Strong compatibility and \mathcal{L} -topologies	17
7	Miscellaneous	19
7.1	Some examples	19
7.2	CANTOR-BENDIXSON-topology	20

1 Introduction

Topological methods are useful in the theory of ω -languages in connection with proving hierarchy results (e.g. [Tho90, Sta97, PP04]). To this end one considers, for a finite alphabet X , the set of all infinite words (ω -words) over X as the infinite product space of the discrete space X . This topology is also known as the CANTOR topology. Infinite words may also be viewed as limits of (infinite) increasing families of finite words w.r.t. the

prefix ordering. Thus it seems to be providing to include both into the same space. One attempt into this direction was done by Boasson and Nivat [BN80]. Redziejowski [Red86] observed that the limit considered in [BN80] is different from that one used in the theory of ω -automata. Therefore he proposed another topology including finite and infinite words into one space. Recently, in [CJS09] possibilities to extend right topologies generated by partial orders on X^* to topologies on X^ω or $X^* \cup X^\omega$ were investigated.

Each of these concepts seems to have several drawbacks when considering topology in connection with acceptance of ω -words, the first approach yields a trivial topological structure for finite words whereas the other ones for infinite words give topologies others than the topology of the CANTOR space:

The topology considered by Boasson and Nivat is closely related to the product or CANTOR topology of X^ω , on the one hand its restriction to X^ω is the CANTOR topology and, on the other hand the whole space $X^* \cup X^\omega$ is a homeomorphic to a closed subset of the CANTOR space $(X \cup \{\perp\})^\omega$ where $\perp \notin X$ is a new letter. However, all finite word languages are open in this topology, in particular, each finite word is an isolated point in this topology.

Redziejowski's topology has all sets consisting of only infinite words (ω -languages) as closed sets, thus providing no information on the complexity of acceptance by topological means.

Finally, the results of [CJS09] show that unless one uses the prefix order \sqsubseteq the limit process extending the topology on X^* yields ambiguous results, more precisely, the limit set may contain more than one element and the topology induced on the space of infinite words may not coincide with the CANTOR topology.

In this paper we investigate the possibility to link topologies on the countable set X^* to the standard topology [Tho90, Sta97, PP04] of the CANTOR space X^ω via a mapping preserving closedness and openness of sets.

As a mapping we use the δ -limit introduced in [Dav64]. Because of the countability of X^* we cannot expect to obtain a full topological correspondence between the spaces X^* and X^ω via any mapping. The δ -limit does not go beyond the class \mathbf{G}_δ of the BOREL hierarchy. For the prefix topology, it provides a correspondence between the open and closed

sets in X^* and the CANTOR space, respectively, and furthermore, of the (σ, δ) -subsets of X^* as introduced in [Sta87] and subsets of X^ω being simultaneously of type \mathbf{F}_σ and \mathbf{G}_δ (see [Sta97, Section 2.4]).

Among the topologies on X^* we consider here are the topologies induced by the $\text{Anf}_\mathcal{L}$ -operators introduced in [PU79, Pro80, Pro83]. Particular cases of these topologies are the topologies induced by the centers [PU79, BN80, BN81] and supercenters [SN86] of languages. It turns out that both topologies play a special rôle: the center-topology being the coarsest one having all finite sets closed, and the supercenter-topology being the finest one for which the prefix set $\mathbf{pref}(F)$ of every closed subset $F \subseteq X^\omega$ is the smallest closed set in X^* corresponding to F via the δ -limit.

The paper is organised as follows. After the notation used we introduce some topological background emphasising properties of the derived set operator. Then we deal with several topologies for finite and infinite words. Section 4 describes the linking of topologies for finite words with the CANTOR topology for ω -words. A fundamental property for a topology of finite words is the compatibility with the topology of infinite words. The class of \mathcal{L} -topologies introduced by Prodinger and Urbanek [PU79, Pro80] is the topic of Section 5. In the subsequent part we deal with topologies fulfilling a strengthened compatibility condition. Some of these results were announced in [Sta10]. Finally we show some limitations of \mathcal{L} -topologies in respect to compatibility with the CANTOR topology.

2 Preliminaries

2.1 Notation

We introduce the notation used throughout the paper. By $\mathbb{N} = \{0, 1, 2, \dots\}$ we denote the set of natural numbers. Let X be an alphabet of cardinality $|X| \geq 2$, $a, b \in X, a \neq b$. By X^* we denote the set (monoid) of words on X , including the *empty word* e , and X^ω is the set of infinite sequences (ω -words) over X . For $w \in X^*$ and $\eta \in X^* \cup X^\omega$ let $w \cdot \eta$ be their *concatenation*. This concatenation product extends in an obvious way to subsets $W \subseteq X^*$ and $B \subseteq X^* \cup X^\omega$. For a language W let $W^* := \bigcup_{i \in \mathbb{N}} W^i$ be the *submonoid* of X^* generated by W , and by $W^\omega := \{w_1 \cdots w_i \cdots : w_i \in W \setminus \{e\}\}$

we denote the set of infinite strings formed by concatenating words in W . Furthermore $|w|$ is the *length* of the word $w \in X^*$ and $\mathbf{pref}(B)$ is the set of all finite prefixes of strings in $B \subseteq X^* \cup X^\omega$. We shall abbreviate $w \in \mathbf{pref}(\eta)$ ($\eta \in X^* \cup X^\omega$) by $w \sqsubseteq \eta$.

Further we denote by $B/w := \{\eta : w \cdot \eta \in B\}$ the *left derivative (left quotient)* of the set $B \subseteq X^* \cup X^\omega$ generated by the word w .

In the theory of ω -automata, ω -words are introduced as upper bounds of infinite chains of words ordered by the prefix relation “ \sqsubseteq ”; this is reflected by the following limit operation (see [CJS09, Dav64, Red86, Sta87, Sta97]).

The δ -limit of a language $W \subseteq X^*$ is defined as

$$W^\delta := \{\xi : \xi \in X^\omega \wedge |\mathbf{pref}(\xi) \cap W| = \aleph_0\}. \quad (1)$$

2.2 General topology

Usually, a *topology* $\mathcal{T} = (\mathcal{X}, \mathcal{O})$ on a set (space) \mathcal{X} is given by a family of open sets $\mathcal{O} \subseteq 2^{\mathcal{X}}$. Here \mathcal{O} is a family of subsets of \mathcal{X} containing \mathcal{X} and closed under arbitrary (including empty) union and finite intersection.

Following Kuratowski (cf. [Kur66, RS63]) one can also define a topology via a closure operator. A mapping $\alpha : 2^{\mathcal{X}} \rightarrow 2^{\mathcal{X}}$ is called a *topological closure operator* provided it satisfies the following conditions.

$$\alpha(\emptyset) = \emptyset \quad (2)$$

$$\alpha(M) \supseteq M \quad (3)$$

$$\alpha(M_1 \cup M_2) = \alpha(M_1) \cup \alpha(M_2), \text{ and} \quad (4)$$

$$\alpha(\alpha(M)) = \alpha(M) \quad (5)$$

In view of Eq. (3) the identity in Eq. (5) can be replaced by $\alpha(\alpha(M)) \subseteq \alpha(M)$.

By \mathcal{T}_α we denote the topology $\mathcal{T}_\alpha = (\mathcal{X}, \mathcal{O}_\alpha)$ where $\mathcal{O}_\alpha := \{\mathcal{X} \setminus \alpha(M) : M \subseteq \mathcal{X}\}$.

A topology $\mathcal{T}_1 = (\mathcal{X}, \mathcal{O}_1)$ is *finer*¹ than a topology $\mathcal{T}_2 = (\mathcal{X}, \mathcal{O}_2)$ if $\mathcal{O}_2 \subseteq \mathcal{O}_1$, that is, if every \mathcal{T}_2 -open set is also \mathcal{T}_1 -open. This is equivalent to the fact that it holds

$$\alpha_1(M) \subseteq \alpha_2(M) \text{ for all } M \in \mathcal{X} \quad (6)$$

¹This includes the case that \mathcal{T}_1 and \mathcal{T}_2 coincide.

where $\alpha_i, i = 1, 2$ is the closure generating the topology \mathcal{T}_i .

A closure operator may also be obtained via the CANTOR-BENDIXSON *derived set* (set of *accumulation points*) M^d of $M \subseteq \mathcal{X}$ [Kur66, Ch. 1, § 9, III].

The derived set operator d has to satisfy the following conditions.

$$\emptyset^d = \emptyset \quad (7)$$

$$(M_1 \cup M_2)^d = M_1^d \cup M_2^d, \text{ and} \quad (8)$$

$$(M^d)^d \subseteq M^d \quad (9)$$

In particular, every closure α is also a derived set operator.

It is readily seen that $\alpha^d(M) := M \cup M^d$ is a topological closure operator and in view of Eq. (9) M^d is closed in \mathcal{T}_{α^d} , in general, a set M is closed in \mathcal{T}_{α^d} if and only if $M^d \subseteq M$. Via α^d , a derived set operator d defines a topology on \mathcal{X} .

The following properties hold (cf. also [Kur66, Ch. 1, § 7, IV and § 9, III]).

Property 1 *Every M' , where $M^d \subseteq M' \subseteq M \cup M^d$, is closed in \mathcal{T}_{α^d} .*

Property 2 *Let $\gamma : 2^{\mathcal{X}} \rightarrow 2^{\mathcal{X}}$ be a derived set or closure operator. Then*

$$G \cap \gamma(M) = G \cap \gamma(G \cap M)$$

if G is open in the topology defined by γ .

For the sake of completeness we add a proof.

Proof. The inclusion \supseteq is obvious.

From Eq. (8) we have $\gamma(M) = \gamma(M \cap G) \cup \gamma(M \setminus G) \subseteq \gamma(M \cap G) \cup \gamma(\mathcal{X} \setminus G)$. Since $\mathcal{X} \setminus G$ is closed, $\gamma(\mathcal{X} \setminus G) \subseteq \mathcal{X} \setminus G$, and we have $\gamma(M) \subseteq \gamma(M \cap G) \cup (\mathcal{X} \setminus G)$ whence the reverse inclusion follows via intersection with the set G . \square

As usual countable unions of closed sets are called \mathbf{F}_σ -sets and countable intersections of open sets are called \mathbf{G}_δ -sets.

3 Topologies for Words and ω -words

In this section we present some known topologies for finite and infinite words.

3.1 The CANTOR topology on X^ω

The first one is the widely investigated CANTOR topology.

We consider the set X^ω as a metric space (CANTOR space) (X^ω, ρ) of all ω -words over the alphabet X where the metric ρ is defined as follows.

$$\rho(\xi, \eta) := \inf\{|X|^{-|w|} : w \sqsubset \xi \wedge w \sqsubset \eta\} \quad (10)$$

Since X is finite, this space is compact, and $\mathcal{C}(F) := \{\xi : \mathbf{pref}(\xi) \subseteq \mathbf{pref}(F)\}$ is the *closure* of the set F (smallest closed subset containing F) in (X^ω, ρ) . Thus

$$\mathbf{pref}(\mathcal{C}(F)) = \mathbf{pref}(F) \text{ and } \mathcal{C}(F) = \mathbf{pref}(F)^\delta. \quad (11)$$

The topology can be defined by the metric ρ as in Eq. (10) or, alternatively, by letting $\mathcal{O}_C := \{W \cdot X^\omega : W \subseteq X^*\}$ be the set of open subsets of X^ω . Then a subset $F \subseteq X^\omega$ is closed if and only if $\mathbf{pref}(\xi) \subseteq \mathbf{pref}(F)$ implies $\xi \in F$.

As (X^ω, ρ) is a metric space the CANTOR-BENDIXSON *derived set* F^d of a subset $F \subseteq X^\omega$ can be described as

$$F^d = \{\xi : \mathbf{pref}(\xi) \subseteq \{w : w \cdot X^\omega \cap F \text{ is infinite}\}\} \quad (12)$$

It holds $\mathcal{C}(F) = F \cup F^d$.

We conclude this part with a relation between the δ -limit and \mathbf{G}_δ -sets in CANTOR space.

Proposition 3 ([Dav64]) *A subset $F \subseteq X^\omega$ is a \mathbf{G}_δ -set if and only if there is a language $W \subseteq X^*$ such that $F = W^\delta$.*

3.2 The prefix topology \mathcal{T}_p on X^*

A well-known topology on X^* which resembles the product or CANTOR topology is the prefix topology \mathcal{T}_p on X^* , that is, the right topology on X^* derived from the prefix order \sqsubseteq on X^* .

The open subsets in \mathcal{T}_p are of the form $W \cdot X^*$ where $W \subseteq X^*$. As it is easily verified the closure operator defining this topology is the initial word operator \mathbf{pref} assigning to each language $W \subseteq X^*$ its set of prefixes $\mathbf{pref}(W) := \{w : \exists v (v \in W \wedge w \sqsubseteq v)\}$.

It has the following property.

Lemma 4 *The prefix topology \mathcal{T}_p is the coarsest topology having all subsets $\mathbf{pref}(F) \subseteq X^*$, $F \subseteq X^\omega$, closed.*

Proof. This follows from the identity $\mathbf{pref}(W) = \mathbf{pref}(W \cdot a^\omega) \cap \mathbf{pref}(W \cdot b^\omega)$ where $a, b \in X$, $a \neq b$. \square

In this topology, however, not every finite set $W \subseteq X^*$ is closed. It holds only the so-called T_0 -condition: For every pair $w, v \in X^*$, $w \neq v$, there is an open set $W \subseteq X^*$ which contains exactly one of the words w or v .

3.3 Topologies on $X^* \cup X^\omega$

In this section we consider several topologies on the space of finite and infinite words $X^* \cup X^\omega$ and its restrictions to the sets X^ω and X^* , respectively.

The first one resembles the Scott topology on $X^* \cup X^\omega$ (cf. [Smy92]). Its open sets are of the form $W \cdot (X^* \cup X^\omega)$. The restriction to X^ω is the CANTOR topology and the restriction to X^* is the prefix topology \mathcal{T}_p .

The topology considered by Boasson and Nivat [BN80] is closely related to the product or CANTOR topology of X^ω , on the one hand its restriction to X^ω is the CANTOR topology and, on the other hand the whole space $X^* \cup X^\omega$ is a homeomorphic to a closed subset of the CANTOR space $(X \cup \{\perp\})^\omega$ where $\perp \notin X$ is a new letter. However, all finite word languages are open in this topology, in particular, each finite word is an isolated point in this topology. Thus its restriction to X^* is the (trivial) discrete topology.

Redziejowski [Red86] observed that the limit considered in [BN80] is different from the one used in the theory of ω -automata. Therefore he proposed another topology on the space of finite and infinite words. Here the closure of a subset $W \cup F \subseteq X^* \cup X^\omega$, $W \subseteq X^*$, $F \subseteq X^\omega$, can be described via the δ -limit as (cf. [Red86, Property 4.4 (1)])

$$\text{cl}(W \cup F) = W \cup W^\delta \cup F.$$

Then subsets of the form $F \subseteq X^\omega$ and $W \cup X^\omega$, $W \subseteq X^*$, are closed, and, consequently, the restrictions to X^ω and X^* , respectively, are the (trivial) discrete topologies.

4 Linking Topologies on X^* to CANTOR Topology

The open sets in \mathcal{T}_p resemble those in CANTOR space. One easily observes, that due to the identity $(W \cdot X^*)^\delta = W \cdot X^\omega$, Eq. (11) and Lemma 4 there is a close correspondence between open (closed) subsets in the prefix topology on X^* and CANTOR topology, respectively.

4.1 Compatibility of topologies

The connection between the prefix topology on X^* and the CANTOR topology on X^ω via the δ -limit is shown in [Sta97, Section 2.4]. This connection fulfils the following property.

Definition 1 (Compatibility) A topology $\mathcal{T} = (X^*, \mathcal{O})$ is *compatible* with the CANTOR topology of X^ω provided

1. W^δ is closed (open) if $W \subseteq X^*$ is closed (open, respectively) in (X^*, \mathcal{O}) .
2. If $F \subseteq X^\omega$ is closed (open) in CANTOR space then $F = W^\delta$ for some $W \subseteq X^*$ closed (open, respectively) in (X^*, \mathcal{O}) , and

Therefore we consider topologies on X^* which are linked via our δ -limit to the CANTOR topology on X^ω . Then every language $W \subseteq X^*$ has as its image the \mathbf{G}_δ -set $W^\delta \subseteq X^\omega$. (In fact, because of the different cardinalities of the spaces X^* and X^ω , we cannot expect to obtain every subset of X^ω as an image.)

Definition 1 requires that the image of every open (closed) language $W \subseteq X^*$ is also open (closed), and every open (closed) ω -language $F \subseteq X^\omega$ is the image of an appropriately chosen open (closed) language.

Lemma 5 *Let a topology \mathcal{T}_α be compatible with the CANTOR topology. Then the closure $\alpha: 2^{X^*} \rightarrow 2^{X^*}$ satisfies the following inclusions*

$$\mathbf{pref}(W^\delta) \subseteq \mathbf{pref}(\alpha(W)^\delta) \subseteq \mathbf{pref}(\alpha(W)) \text{ and} \quad (13)$$

$$\mathcal{C}(W^\delta) \subseteq \alpha(W)^\delta, \text{ for all } W \subseteq X^*. \quad (14)$$

Proof. The first inclusion follows from the facts that $W \subseteq \alpha(W)$ and $\mathbf{pref}(V^\delta) \subseteq \mathbf{pref}(V)$, and the second one from the fact that $\alpha(W)^\delta$ is a closed set containing W^δ . \square

Moreover, Conditions 1. and 2. of Definition 1 are inherited in the following way.

Lemma 6 *Let $\mathcal{T} = (X^*, \mathcal{O})$ be a topology compatible with the CANTOR topology.*

1. *If the topology $\mathcal{T}_1 = (X^*, \mathcal{O}_1)$ is coarser than \mathcal{T} then the δ -image W^δ of every open (closed) subset $W \subseteq X^*$ is open (closed, respectively) in X^ω .*
2. *If the topology $\mathcal{T}_2 = (X^*, \mathcal{O}_2)$ is finer than \mathcal{T} then every open (closed) subset $F \subseteq X^\omega$ has a open (closed, respectively) δ -preimage $W \subseteq X^*$.*

Lemma 6.2 has the following special instance.

Corollary 7 *If a topology \mathcal{T} on X^* is finer than the prefix topology \mathcal{T}_p then every closed (open) subset $F \subseteq X^\omega$ ($W \cdot X^\omega$) has the closed (open) δ -preimage $\mathbf{pref}(F)$ ($W \cdot X^*$).*

4.2 (σ, δ) -subsets of X^*

As Proposition 3 shows, the δ -limit can map languages only to ω -languages in the BOREL-class \mathbf{G}_δ , it is interesting to observe that we can also characterise those \mathbf{G}_δ -sets which are simultaneously \mathbf{F}_σ -sets via their δ -preimages (see [Sta87]). These subsets of X^* show also some special properties w.r.t. topologies on X^* .

We start with some easily verified properties of the δ -limit defined in Eq. (1).

$$(W \cup V)^\delta = W^\delta \cup V^\delta \quad (15)$$

$$\mathbf{pref}(\mathbf{pref}(W^\delta)^\delta) = \mathbf{pref}(W^\delta) \subseteq \mathbf{pref}(W) \quad (16)$$

The second identity follows from applying Eq. (11) to $F = W^\delta$.

Definition 2 A subset $W \subseteq X^*$ is referred to as a (σ, δ) -subset of X^* provided for every $\xi \in X^\omega$ one of the sets $\mathbf{pref}(\xi) \cap W$ or $\mathbf{pref}(\xi) \setminus W$ is finite.²

Then we have the announced connection to \mathbf{F}_σ -sets in CANTOR space.

Lemma 8 ([Sta87, Lemma 12]) *A subset $F \subseteq X^\omega$ is simultaneously an \mathbf{F}_σ - and a \mathbf{G}_δ -set in (X^ω, ρ) if and only if there is a (σ, δ) -subset $W \subseteq X^*$ such that $F = W^\delta$.*

²There are languages $W \subseteq X^*$ such that both sets $\mathbf{pref}(\xi) \cap W$ and $\mathbf{pref}(\xi) \setminus W$ are infinite.

Examples of (σ, δ) -subsets of X^* are languages of the form $\mathbf{pref}(W)$, $W \cdot X^*$, and $W \subseteq X^*$ such that $W^\delta = \emptyset$.

Definition 2 is equivalent to the following one

$$W \text{ is a } (\sigma, \delta)\text{-subset of } X^* \text{ if and only if } (X \setminus W)^\delta = X^\omega \setminus W^\delta. \quad (17)$$

We have also the following equivalent property.

Lemma 9 *A subset $W \subseteq X^*$ is a (σ, δ) -subset of X^* if and only if for every $V \subseteq X^*$ one of the identities $(V \cap W)^\delta = V^\delta \cap W^\delta$ or $(V \setminus W)^\delta = V^\delta \setminus W^\delta$ is fulfilled.*

Proof. The inclusion $(V \cap W)^\delta \subseteq V^\delta \cap W^\delta$ is obvious.

If W is not a (σ, δ) -subset of X^* then there is a $\xi \in W^\delta$ such that $V := \mathbf{pref}(\xi) \setminus W$ is infinite. Then $V^\delta = \{\xi\}$ and, consequently, $V^\delta \cap W^\delta = \{\xi\}$ whereas $(V \cap W)^\delta = \emptyset$.

Let W be a (σ, δ) -subset of X^* , and let $\xi \in W^\delta \cap V^\delta$. Then $\xi \in W^\delta$ implies that $\mathbf{pref}(\xi) \setminus W$ is finite. Since $\mathbf{pref}(\xi) \cap V$ is infinite, $\mathbf{pref}(\xi) \cap V \cap W$ is infinite, too.

The other assertion follows if we replace W by $X^* \setminus W$. \square

From Eq. (11) we have $W^\delta \subseteq \mathcal{C}(W^\delta) = \mathbf{pref}(W^\delta)^\delta$. Thus Lemma 9 and Eq. (15) imply the following.

$$(W \cap \mathbf{pref}(W^\delta))^\delta = W^\delta \text{ and } (W \setminus \mathbf{pref}(W^\delta))^\delta = \emptyset. \quad (18)$$

Corollary 10 ([Sta87, Proposition 14]) *The class of all (σ, δ) -subsets of X^* is a Boolean algebra.*

We finish this section with a sufficient condition for the compatibility of topologies $\mathcal{T} = (X^*, \mathcal{O})$ which is immediate from Eq. (17).

Lemma 11 *Let $\mathcal{T} = (X^*, \mathcal{O})$ be a topology on X^* such that every open set $W \in \mathcal{O}$ is a (σ, δ) -subset of X^* . Then \mathcal{T} is compatible with the CANTOR topology if W^δ is open for every $W \in \mathcal{O}$ and every open subset $E \subseteq X^\omega$ has an open δ -preimage $V \in \mathcal{O}$.*

5 \mathcal{L} -topologies

In this section we investigate a method for defining a class of topologies on X^* which are finer than the prefix topology \mathcal{T}_p .

5.1 The operator Anf

This class can be defined via the Anf-operator introduced by Proding and Urbanek [PU79, Pro80]. They defined a generalisation of the initial word operator **pref** as follows. Let $\mathcal{L} \subseteq 2^{X^*}$ be a family of languages, $W \subseteq X^*$, and define

$$\text{Anf}_{\mathcal{L}}(W) := \{w : w \in X^* \wedge W/w \in \mathcal{L}\} \quad (19)$$

The operator $\text{Anf}_{\mathcal{L}}$ has the following properties (see [PU79, Pro80, RS08]):

$$\text{Anf}_{\mathcal{L}}(W) \subseteq \text{Anf}_{\mathcal{M}}(W) \text{ if and only if } \mathcal{L} \subseteq \mathcal{M} \quad (20)$$

Property 12 1. $\text{Anf}_{\mathcal{L}}(W/w) = \text{Anf}_{\mathcal{L}}(W)/w$

2. $\text{Anf}_{\mathcal{L}}$ is monotone if and only if $W \in \mathcal{L}$ and $W \subseteq V$ imply $V \in \mathcal{L}$.

3. $\text{Anf}_{\mathcal{L}}$ is \cup -stable if and only if $\text{Anf}_{\mathcal{L}}$ is monotone and $W \cup V \in \mathcal{L}$ implies $W \in \mathcal{L}$ or $V \in \mathcal{L}$.

Property 13 The following conditions are equivalent.

1. $\emptyset \notin \mathcal{L}$
2. $\text{Anf}_{\mathcal{L}}(\emptyset) = \emptyset$, and
3. $\forall W (\text{Anf}_{\mathcal{L}}(W) \subseteq \mathbf{pref}(W))$

Proof. 1 \rightarrow 3. Let $\emptyset \notin \mathcal{L}$ and $v \notin \mathbf{pref}(W)$. Then $W/v = \emptyset \notin \mathcal{L}$ whence $v \notin \text{Anf}_{\mathcal{L}}(W)$.

3 \rightarrow 2. is obvious

2 \rightarrow 1. We have $e \in \text{Anf}_{\mathcal{L}}(W)$ if and only if $W \in \mathcal{L}$. Thus $e \notin \text{Anf}_{\mathcal{L}}(\emptyset) = \emptyset$ implies $\emptyset \notin \mathcal{L}$. \square

Next we give some relations to the prefix-operator $\mathbf{pref} = \text{Anf}_{\mathcal{L}_p}$.

Proposition 14 Let $\mathcal{L} \subseteq 2^{X^*}$. The following conditions are equivalent.

1. $\forall W (W \subseteq X^* \rightarrow (\text{Anf}_{\mathcal{L}}(W) \neq \emptyset \rightarrow W \in \mathcal{L}))$
2. $\forall W \forall v (W \subseteq X^* \wedge v \in X^* \rightarrow (W/v \in \mathcal{L} \rightarrow W \in \mathcal{L}))$
3. $\forall W (W \subseteq X^* \rightarrow \mathbf{pref}(\text{Anf}_{\mathcal{L}}(W)) = \text{Anf}_{\mathcal{L}}(W))$

Proof. The equivalence of Items 1 and 2 is Theorem 4.18 of [PU79].

3 \rightarrow 1. If $W/v \in \mathcal{L}$ then $v \in \text{Anf}_{\mathcal{L}}(W)$. Now 3. implies $e \in \mathbf{pref}(\text{Anf}_{\mathcal{L}}(W)) \subseteq \text{Anf}_{\mathcal{L}}(W)$ which in turn shows $W \in \mathcal{L}$.

2 \rightarrow 3. Let $w \in \mathbf{pref}(\text{Anf}_{\mathcal{L}}(W))$. Then there is a $v \in X^*$ such that $w \cdot v \in \text{Anf}_{\mathcal{L}}(W)$, that is, $W/w \cdot v = (W/w)/v \in \mathcal{L}$. Now 2. yields $W/w \in \mathcal{L}$ which in turn shows $w \in \text{Anf}_{\mathcal{L}}(W)$. \square

We continue with the requirements under which the operator $\text{Anf}_{\mathcal{L}}$ satisfies the conditions of Eqs. (9) of a derived set operator.

Lemma 15 *It holds $\forall W (\text{Anf}_{\mathcal{L}}(\text{Anf}_{\mathcal{L}}(W)) \subseteq \text{Anf}_{\mathcal{L}}(W))$ if and only if $\forall W (\text{Anf}_{\mathcal{L}}(W) \in \mathcal{L} \rightarrow W \in \mathcal{L})$.*

Proof. Let $\text{Anf}_{\mathcal{L}}(W) \in \mathcal{L}$. Consequently, $e \in \text{Anf}_{\mathcal{L}}(\text{Anf}_{\mathcal{L}}(W))$. Then the inclusion $\text{Anf}_{\mathcal{L}}(\text{Anf}_{\mathcal{L}}(W)) \subseteq \text{Anf}_{\mathcal{L}}(W)$ implies $e \in \text{Anf}_{\mathcal{L}}(W)$. Thus $W \in \mathcal{L}$.

Conversely, let $\text{Anf}_{\mathcal{L}}(W) \in \mathcal{L}$ imply $W \in \mathcal{L}$, and let $v \in \text{Anf}_{\mathcal{L}}(\text{Anf}_{\mathcal{L}}(W))$. Then $\text{Anf}_{\mathcal{L}}(W)/v \in \mathcal{L}$. Since $\text{Anf}_{\mathcal{L}}(W)/v = \text{Anf}_{\mathcal{L}}(W/v)$, we have $\text{Anf}_{\mathcal{L}}(W/v) \in \mathcal{L}$ and, consequently, $W/v \in \mathcal{L}$, that is $v \in \text{Anf}_{\mathcal{L}}(W)$. \square

5.2 \mathcal{L} -topologies

Under several conditions an $\text{Anf}_{\mathcal{L}}$ -operator has the properties of a derived set operator on X^* . The following theorem is an analogue to Theorem 2.3 of [Pro80] for derived set operators.

Theorem 16 *A mapping $\text{Anf}_{\mathcal{L}}$ is a derived set operator on X^* if and only if the following conditions are satisfied.*

1. $\emptyset \notin \mathcal{L}$,
2. $W \in \mathcal{L}$ and $W \subseteq V$ imply $V \in \mathcal{L}$, and $W \cup V \in \mathcal{L}$ implies $W \in \mathcal{L}$ or $V \in \mathcal{L}$, and
3. $\text{Anf}_{\mathcal{L}}(W) \in \mathcal{L}$ implies $W \in \mathcal{L}$.

Proof. 1. First $\emptyset \in \mathcal{L}$ if and only if $\text{Anf}_{\mathcal{L}}(\emptyset) = \emptyset$. Thus Eq. (7) holds.

2. Theorem 4.13 of [PU79] shows that this is equivalent to $\text{Anf}_{\mathcal{L}}(V \cup W) = \text{Anf}_{\mathcal{L}}(V) \cup \text{Anf}_{\mathcal{L}}(W)$ which in turn is Eq. (8).

3. Lemma 15 shows that Item 3 is equivalent to Eq. (9). \square

Simple examples of \mathcal{L} -topologies are the prefix topology \mathcal{T}_p and the discrete topology $\mathcal{D} = (X^*, 2^{X^*})$. Here one can choose $\mathcal{L}_p = \{W : W \neq \emptyset\}$, and

$\mathcal{L}_{\mathcal{D}} = \emptyset$, respectively. This yields $\text{Anf}_{\mathcal{L}_p} = \mathbf{pref}$ and $\text{Anf}_{\mathcal{L}_{\mathcal{D}}}(W) = \emptyset$, for all $W \subseteq X^*$.

Utilising Property 13.3 and Proposition 14.3 we obtain the following sufficient condition for $\text{Anf}_{\mathcal{L}}$ to be a derived set operator.

Corollary 17 *A mapping $\text{Anf}_{\mathcal{L}}$ is a derived set operator on X^* if the following conditions are satisfied.*

1. $\emptyset \notin \mathcal{L}$,
2. $W \in \mathcal{L}$ and $W \subseteq V$ imply $V \in \mathcal{L}$, and $W \cup V \in \mathcal{L}$ implies $W \in \mathcal{L}$ or $V \in \mathcal{L}$, and
3. If $W/w \in \mathcal{L}$ then $W \in \mathcal{L}$.

Since every \mathcal{L} -topology is finer than the prefix topology \mathcal{T}_p , Lemma 6.2 and Corollary 7 yield the following.

Proposition 18 An \mathcal{L} -topology is *compatible* with the CANTOR topology on X^ω if and only if for every $W \subseteq X^*$, the set $\alpha_{\mathcal{L}}(W)^\delta$ is closed and the set $(X^* \setminus \alpha_{\mathcal{L}}(W))^\delta$ is open in CANTOR space.

Finally, we mention that \mathcal{L} -spaces can be characterised also via their open sets.

Theorem 19 ([Pro80, Theorem 2.16]) *A topological space $\mathcal{T} = (X^*, \mathcal{O})$ is an \mathcal{L} -space if and only if $W \in \mathcal{O}$ implies $w \cdot W, W/w \in \mathcal{O}$ for $w \in X^*$.*

6 Strongly Compatible Topologies on X^*

In this section we consider topologies on X^* compatible with the CANTOR topology. Several of them can be defined using the apparatus introduced in the preceding section.

6.1 Center and Supercenter topologies

Special $\text{Anf}_{\mathcal{L}}$ -operators were considered in connection with language-theoretic questions. These were referred to as centers [BN81] or supercenters [SN86], respectively, of languages.

Definition 3 (Center)

$$\text{center}(W) := \text{Anf}_{\mathcal{L}_c}(W), \text{ where } \mathcal{L}_c = \{V : V \subseteq X^* \text{ is infinite}\}$$
Definition 4 (Supercenter)

$$\text{sctr}(W) := \text{Anf}_{\mathcal{L}_{sc}}(W), \text{ where } \mathcal{L}_{sc} = \{V : V \subseteq X^* \wedge V^\delta \neq \emptyset\}$$

Definition 4 is equivalent to $\text{sctr}(W) = \mathbf{pref}(W^\delta) = \mathbf{pref}(\mathcal{C}(W^\delta))$.

Since the alphabet X is finite, König's infinity lemma shows that $W \subseteq X^*$ is infinite if and only if $\mathbf{pref}(W)$ is infinite which is equivalent to $\mathbf{pref}(W)^\delta \neq \emptyset$. Thus we have

$$\text{center}(W) = \text{sctr}(\mathbf{pref}(W)) = \mathbf{pref}(\mathbf{pref}(W)^\delta). \quad (21)$$

Both families \mathcal{L}_c and \mathcal{L}_{sc} satisfy the conditions of Corollary 17. So center and sctr are derived set operators and define topologies \mathcal{T}_c and \mathcal{T}_{sc} , respectively, on X^* . As \mathcal{L} -topologies the center or supercenter topologies are finer than the prefix topology \mathcal{T}_p . Moreover, Eq. (20) shows that \mathcal{T}_{sc} is finer than \mathcal{T}_c .

In \mathcal{T}_{sc} every W with $W^\delta = \emptyset$ is closed. The language $a^* \cdot b$ is infinite, $(a^* \cdot b)^\delta = \emptyset$ and $\text{center}(a^* \cdot b) = a^* \not\subseteq a^* \cdot b$. Therefore it is not closed in \mathcal{T}_c . This shows that \mathcal{T}_{sc} does not coincide with \mathcal{T}_c .

The prefix topology has the property that not every finite subset of X^* is closed. The center topology proves to be the coarsest topology refining \mathcal{T}_p and having all finite sets closed. To this end we show the following.

Theorem 20 *If $\alpha: 2^{X^*} \rightarrow 2^{X^*}$ is a topological closure such that all $\mathbf{pref}(F), F \subseteq X^\omega$, and all finite sets are closed then*

$$\alpha(W \cup \text{center}(W)) = W \cup \text{center}(W).$$

Proof. If all $\mathbf{pref}(F), F \subseteq X^\omega$, are α -closed, then according to Lemma 4 $\alpha(W) \subseteq \mathbf{pref}(W)$, for all $W \subseteq X^*$, and all $W \cdot X^*$ are α -open.

Eq. (21) shows that every $\text{center}(W)$ is α -closed. Now the inclusion $\alpha(W \cup \text{center}(W)) \supseteq W \cup \text{center}(W)$ follows from $\alpha(V) \supseteq V$.

It suffices to show $\alpha(W) \setminus \text{center}(W) \subseteq W$.

Let $u \in \alpha(W) \setminus \text{center}(W) \subseteq \mathbf{pref}(W) \setminus \text{center}(W)$. Then $W \cap u \cdot X^*$ is finite. Consequently, $\alpha(W \cap u \cdot X^*) = W \cap u \cdot X^*$. Since $u \cdot X^*$ is α -open, Property 2 shows $\alpha(W) \cap u \cdot X^* = \alpha(W \cap u \cdot X^*) \cap u \cdot X^* = W \cap u \cdot X^*$ whence $u \in W$. \square

Corollary 21 *The topology \mathcal{T}_c is the coarsest topology having all subsets $\mathbf{pref}(F) \subseteq X^*$ and all finite subsets closed.*

Proof. This follows from Eq. (6) and Theorem 20. \square

The next lemma shows a connection between the supercenter topology and (σ, δ) -subsets of X^* .

Lemma 22 *If a topology \mathcal{T} on X^* is coarser than the supercenter topology \mathcal{T}_{sc} then every open or closed set is a (σ, δ) -subsets of X^* .*

Proof. If \mathcal{T} is coarser than \mathcal{T}_{sc} , then every $V \subseteq X^*$ closed in \mathcal{T} is also closed in \mathcal{T}_{sc} , that is, is of the form $V = W \cup \mathbf{pref}(W^\delta) = (W \setminus \mathbf{pref}(W^\delta)) \cup \mathbf{pref}(W^\delta)$. Along with all prefix-closed languages, in \mathcal{T}_{sc} all languages W with $\text{sctr}(W) = \mathbf{pref}(W^\delta) = \emptyset$ are closed. Now the assertion follows with Eq. (18) and Corollary 10. \square

The proof of Lemma 22 shows also that every set open in the supercenter topology has the form $W \cdot X^* \setminus V$ where $V^\delta = \emptyset$.

6.2 Strong compatibility - characterisation

In every topology finer than the prefix topology \mathcal{T}_p all languages $\mathbf{pref}(F)$ where $F \subseteq X^\omega$ are closed. Moreover, $\mathbf{pref}(F)^\delta = \mathcal{C}(F)$. In this section we are investigating which topologies have the languages $\mathbf{pref}(F)$ as smallest closed sets yielding $V^\delta = \mathcal{C}(F)$. It turns out that supercenters of languages play a crucial rôle in this respect.

Definition 5 (Strong compatibility) A topology $\mathcal{T} = (X^*, \mathcal{O})$ is *strongly compatible* with the CANTOR topology provided \mathcal{T} is compatible with the CANTOR topology and

$$\forall F (F \subseteq X^\omega \rightarrow \mathbf{pref}(F) = \min_{\subseteq} \{\alpha_{\mathcal{T}}(W) : F \subseteq W^\delta\}) \quad (22)$$

In particular, every $\mathbf{pref}(F)$, $F \subseteq X^\omega$, is closed in \mathcal{T} .

Lemma 23 *A topology \mathcal{T} on X^* is strongly compatible with the CANTOR topology if and only if the corresponding closure operator $\alpha_{\mathcal{T}}$ satisfies $\mathbf{pref}(W) \supseteq \alpha_{\mathcal{T}}(W) \supseteq \mathbf{pref}(W^\delta)$ for all $W \subseteq X^*$.*

Proof. Let \mathcal{T} be strongly compatible with the CANTOR topology. Then every $\mathbf{pref}(F)$ is closed in \mathcal{T} , and Lemma 4 shows that \mathcal{T} refines the prefix topology \mathcal{T}_p .

Moreover, Eq. (22), for $F = W^\delta$, yields $\mathbf{pref}(W^\delta) \subseteq \alpha_{\mathcal{T}}(W)$.

If $\mathbf{pref}(W) \supseteq \alpha_{\mathcal{T}}(W) \supseteq \mathbf{pref}(W^\delta)$ then every $\mathbf{pref}(F)$, $F \subseteq X^\omega$, is closed, and $\mathbf{pref}(F) \in \{\alpha_{\mathcal{T}}(W) : F \subseteq W^\delta\}$, since $F \subseteq \mathbf{pref}(F)^\delta$. Let $\alpha_{\mathcal{T}}(V) \in \{\alpha_{\mathcal{T}}(W) : F \subseteq W^\delta\}$. Then $V^\delta \supseteq F$. Consequently, $\alpha_{\mathcal{T}}(V) \supseteq \mathbf{pref}(V^\delta) \supseteq \mathbf{pref}(F)$.

The compatibility with the CANTOR topology follows from Corollary 7 and Lemma 22. \square

In view of Eq. (6) the inequality of Lemma 23 is equivalent to the condition that the topology \mathcal{T} is finer than the prefix topology \mathcal{T}_p and coarser than the supercenter topology \mathcal{T}_{sc} . Then Lemma 22 implies that open and closed sets are (σ, δ) -subsets of X^* .

6.3 Strong compatibility and \mathcal{L} -topologies

The so far considered strongly compatible topologies $\mathcal{T}_p, \mathcal{T}_c$ and \mathcal{T}_{sc} are \mathcal{L} -topologies. It arises the question whether all strongly compatible topologies are \mathcal{L} -topologies. In view of Lemma 23 this is equivalent to whether all topologies between the prefix and the supercenter topology are \mathcal{L} -topologies.

A further observation is that as a consequence of Lemmata 22 and 23, the closed and the open sets in every topology strongly compatible with the CANTOR topology are always (σ, δ) -subsets of X^* . We also address the question whether all compatible \mathcal{L} -topologies having as open sets (σ, δ) -subsets of X^* are strongly compatible.

In this section we will show that for both instances we find counterexamples. To this end we use the well-known possibility to define topologies via their bases (e.g. [RS63, Ch. I, § 2, 2.1]):

Property 24 *Let X be a set and $\mathcal{B} \subseteq \mathcal{X}$ be closed under intersection. Then for $\mathcal{O} := \{\bigcup_{M \in \mathcal{A}} M : \mathcal{A} \subseteq \mathcal{B}\}$ the pair $\mathcal{T} = (X, \mathcal{O})$ is a topological space with open sets \mathcal{O} .*

Remark. The set \mathcal{B} in Property 24 has the properties of a base of the topological space \mathcal{T} , as it generates all its open sets. In topology, however, it is not required that a base be closed under intersection [Kur66, RS63].

The first example is a topology on X^* which is strongly compatible with the CANTOR topology but not an \mathcal{L} -topology.

Example 1 Let $X = \{a, b\}$ and define $\mathcal{B} := \{w \cdot X^* \setminus V : w \in X^* \wedge V \subseteq a^* \cdot b\}$ and the family of open sets \mathcal{O} as in Property 24. Then $\mathcal{T} = (X^*, \mathcal{O})$ is a topology finer than the prefix topology \mathcal{T}_p and, since $V^\delta = \emptyset$ for $V \subseteq a^* \cdot b$, coarser than \mathcal{T}_{sc} .

Consider $W := b \cdot (X^* \setminus a^* \cdot b)$. Then $b \in W$ and $bb \notin W$. The smallest open set containing the word b is $b \cdot X^* \setminus a \cdot a^* \cdot b$ which contains the word bb . Thus W is not open, and according to Theorem 19 the topology \mathcal{T} is no \mathcal{L} -topology. ✓

Since the topology $\mathcal{T} = (X^*, \mathcal{O})$ of Example 1 is coarser than \mathcal{T}_{sc} , its open sets are (σ, δ) -subsets of X^* . Next we provide an example of a compatible \mathcal{L} -topology having all open sets as (σ, δ) -subsets of X^* but not strongly compatible.

Lemma 25 *There are compatible \mathcal{L} -topologies on X^* such that all its open sets are (σ, δ) -subsets of X^* which are not strongly compatible with the CANTOR topology.*

Proof. We construct an \mathcal{L} -topology $\mathcal{T} = (X^*, \mathcal{O})$ such that every open set $W \in \mathcal{O}$ and hence also every closed set is a (σ, δ) -set. To this end we use Theorem 19 and Property 24.

We let $X = \{a, b\}$ and $\mathcal{B} = \{w \cdot X^* \setminus U \cdot b \cdot a^* : w \in X^* \wedge U \subseteq X^* \text{ finite}\}$ be a base of \mathcal{T} .

Then every open set has the form³

$$W = \bigcup_{i \in M} (w_i \cdot X^* \setminus U_i \cdot b \cdot a^*), \quad M \subseteq \mathbb{N}. \quad (23)$$

First we show that \mathcal{O} is closed under the operations $w \cdot$ and $/w$. Closure under premultiplication with a word is trivial. It remains to show that $(w \cdot X^* \setminus U \cdot b \cdot a^*)/v = (w \cdot X^*/v) \setminus (U \cdot b \cdot a^*)/v$ is a union of sets of the given shape. If $v \notin \mathbf{pref}(w \cdot X^*)$, $(w \cdot X^* \setminus U \cdot b \cdot a^*)/v = \emptyset$. Otherwise, $w \cdot X^*/v = w' \cdot X^*$ for a suffix w' of w . Moreover, observe that

$$(U \cdot b \cdot a^*)/v = \begin{cases} (U/v) \cdot b \cdot a^*, & \text{if } v \notin U \cdot b \cdot a^* \text{ and} \\ (U/v) \cdot b \cdot a^* \cup a^*, & \text{otherwise.} \end{cases}$$

In the former case $(w \cdot X^* \setminus U \cdot b \cdot a^*)/v = w' \cdot X^* \setminus (U/v) \cdot b \cdot a^*$ is of the required form. In the latter case we get $w' \cdot X^* \setminus a^* = \bigcup_{v' \in w' \cdot a^* \cdot b} v' \cdot X^*$ and,

³Observe that the union in Eq. (23) is always an at most countable one.

consequently, $(w \cdot X^* \setminus U \cdot b \cdot a^*)/v = \bigcup_{v' \in w' \cdot a^* \cdot b} (v' \cdot X^* \setminus (U/v) \cdot b \cdot a^*)$ is also in \mathcal{O} .

Next, we show that every open set $W \subseteq X^*$ is a (σ, δ) -subset of X^* , that is, for $\xi \in W^\delta$ we have to show that $\mathbf{pref}(\xi) \setminus W$ is finite. If $\xi \in W^\delta$ in view of Eq. (23) there is an $i \in \mathbb{N}$ such that $\xi \in w_i \cdot X^\omega$. Since U_i is finite, $\xi \notin U_i \cdot b \cdot a^\omega$ implies that $\mathbf{pref}(\xi) \setminus W \subseteq \mathbf{pref}(\xi) \cap U_i \cdot b \cdot a^*$ is also finite.

If $\xi \in U_i \cdot b \cdot a^\omega$ then $\xi = u \cdot b \cdot a^\omega$ for some $u \in U_i$. Since $\mathbf{pref}(\xi) \cap W$ is infinite, there is a $j \in \mathbb{N}$ such that $\mathbf{pref}(\xi) \cap (w_j \cdot X^* \setminus U_j \cdot b \cdot a^*)$, contains a word $|w| \geq |u| + 2$. Thus $u \cdot b \cdot a \sqsubseteq w \sqsubset \xi$.

For $w \in u \cdot b \cdot a^*$ and $w \notin U_j \cdot b \cdot a^*$ it holds $u \notin U_j$. Thus $u \cdot b \cdot a^* \cap U_j \cdot b \cdot a^* = \emptyset$ whence $\mathbf{pref}(\xi) \cap U_j \cdot b \cdot a^* \supseteq \mathbf{pref}(\xi) \setminus W$ is finite.

For the compatibility of $\mathcal{T} = (X^*, \mathcal{O})$ with the CANTOR topology, in view of Lemma 11, it suffices to show that W^δ is open if W is given by Eq. (23). First observe that $W^\delta \subseteq \bigcup_{i \in \mathbb{N}} w_i \cdot X^\omega$ and $w_i \cdot X^\omega \supseteq (W^\delta \cap w_i \cdot X^\omega) \supseteq w_i \cdot X^\omega \setminus U_i \cdot b \cdot a^\omega$. Thus $w_i \cdot X^\omega \setminus W^\delta$ is finite, hence $w_i \cdot X^\omega \cap W^\delta$ is open. Consequently $W^\delta = \bigcup_{i \in \mathbb{N}} (w_i \cdot X^\omega \cap W^\delta)$ is also open.

Finally, the set $b \cdot a^*$ is closed in \mathcal{T} and $(b \cdot a^*)^\delta = \{b \cdot a^\omega\}$ but $\mathbf{pref}(\{b \cdot a^\omega\}) \not\subseteq b \cdot a^*$. Thus $\mathcal{T} = (X^*, \mathcal{O})$ is not strongly compatible with the CANTOR topology. \square

7 Miscellaneous

In this section we first present examples of topologies which are compatible with the CANTOR topology but not strongly compatible. Then we consider \mathcal{L} -topologies related to the CANTOR-BENDIXSON derived set in CANTOR space.

7.1 Some examples

The first one is coarser than the prefix topology, and the second one is a refinement of the first one incomparable with the prefix topology and the center topology. For both topologies the open (and closed) sets are (σ, δ) -sets.

Example 2 Define the topology \mathcal{T}_{2p} by $\mathcal{O}_{2p} := \{W \cdot X^* : W \subseteq (X^2)^*\}$. Then every open set is also a (σ, δ) -subset of X^* .

The set $a \cdot X^* = \{a\} \cup a^2 \cdot X^*$ is not open. Thus \mathcal{T}_{2p} is strictly coarser than \mathcal{T}_p . Since $W \cdot X^\omega = (W \cdot X^* \cap (X^2)^*) \cdot X^\omega = ((W \cdot X^* \cap (X^2)^*) \cdot X^*)^\delta$, Lemma 6.1 shows that \mathcal{T}_{2p} is compatible with the CANTOR topology. ✓

Lemma 26 Let $\mathcal{B}_{2sc} := \{W_i \cdot X^* \setminus V_i : W_i \subseteq (X^2)^* \wedge V_i^\delta = \emptyset\}$ and define \mathcal{O}_{2sc} as in Property 24. Then \mathcal{B}_{2sc} is closed under finite intersection and \mathcal{O}_{2sc} consists solely of (σ, δ) -subsets of X^* .

Proof. Closure under intersection follows from the identities $(W_i \cdot X^* \setminus V_i) \cap (W_j \cdot X^* \setminus V_j) = (W_i \cdot (X^2)^* \cap W_j \cdot (X^2)^*) \cdot X^* \setminus (V_i \cup V_j)$ and $(V_i \cup V_j)^\delta = \emptyset$.

Finally, since $\mathcal{O}_{2sc} \subseteq \mathcal{O}_{sc}$, all open sets are (σ, δ) -subsets of X^* . □

Example 3 Lemma 26 shows that \mathcal{O}_{2sc} is a family of open sets consisting solely of (σ, δ) -sets. Assume $a \cdot X^* = \{a\} \cup a^2 \cdot X^*$ to be open. Then $a \cdot X^* = \bigcup_{i \in I} (W_i \cdot X^* \setminus V_i)$ implies that $e \in W_i \subseteq (X^2)^*$ for some $i \in I$, that is $a \cdot X^* \supseteq X^* \setminus V_i$. This contradicts $a \cdot X^\omega = (a \cdot X^*)^\delta \subseteq (X^* \setminus V_i)^\delta = X^\omega$.

Thus the topology \mathcal{T}_{2sc} is not finer than the prefix topology \mathcal{T}_p . Since every U with $U^\delta = \emptyset$ is closed in \mathcal{T}_{2sc} , in particular, every finite set is closed in \mathcal{T}_{2sc} , the topology is not coarser than \mathcal{T}_c . ✓

7.2 CANTOR-BENDIXSON-topology

In this section we first present two examples of \mathcal{L} -topologies which are strictly finer than the supercenter topology. Thus they are not strongly compatible. The first example is related to the CANTOR-BENDIXSON derived set in CANTOR space and compatible with the CANTOR topology. The second one refers to the CANTOR-BENDIXSON Theorem as it is concerned with condensation points in W^δ , that is, points $\xi \in X^\omega$ for which every $w \cdot X^\omega \cap W^\delta, w \sqsubset \xi$, is uncountable.

Theorem 27 Let $\mathcal{L}_\infty := \{W : W^\delta \text{ is infinite}\}$. Then $\text{Anf}_{\mathcal{L}_\infty}$ is a derived set operator on X^* and the topology defined by $\text{Anf}_{\mathcal{L}_\infty}$ is compatible with the CANTOR topology.

Moreover, $\text{Anf}_{\mathcal{L}_\infty}(W)^\delta = \emptyset$ whenever W^δ is finite.

Proof. First, in view of Proposition 14 and Theorem 16 $\text{Anf}_{\mathcal{L}_\infty}$ is a derived set operator which satisfies $\text{Anf}_{\mathcal{L}_\infty}(W) = \mathbf{pref}(\text{Anf}_{\mathcal{L}_\infty}(W))$. Then $\alpha_{\mathcal{L}_\infty}(W) := W \cup \text{Anf}_{\mathcal{L}_\infty}(W)$ is a closure in X^* .

Next, observe that $w \in \text{Anf}_{\mathcal{L}_\infty}(W)$ is equivalent to W^δ/w is infinite which in turn is equivalent to $W^\delta \cap w \cdot X^\omega$ is infinite. Thus, $\xi \in \text{Anf}_{\mathcal{L}_\infty}(W)^\delta$ if and only if $W^\delta \cap w \cdot X^\omega$ is infinite for all $w \in \mathbf{pref}(\xi)$, that is, in view of Eq. (12) $\text{Anf}_{\mathcal{L}_\infty}(W)^\delta$ is the CANTOR-BENDIXSON derived set of W^δ . This shows that $\alpha_{\mathcal{L}_\infty}(W)^\delta = W^\delta \cup \text{Anf}_{\mathcal{L}_\infty}(W)^\delta = \mathcal{C}(W^\delta)$ is closed.

It remains to prove that $(X^\omega \setminus \alpha_{\mathcal{L}_\infty}(W))^\delta$ is open. To this end we use the fact that $\text{Anf}_{\mathcal{L}_\infty}(W)^\delta$ is the CANTOR-BENDIXSON derived set of W^δ and $\text{Anf}_{\mathcal{L}_\infty}(W) = \mathbf{pref}(\text{Anf}_{\mathcal{L}_\infty}(W))$ is a (σ, δ) -subset of X^* . The latter implies $\text{Anf}_{\mathcal{L}_\infty}(W)^\delta = X^\omega \setminus (X^* \setminus \text{Anf}_{\mathcal{L}_\infty}(W))^\delta$. Then $\text{Anf}_{\mathcal{L}_\infty}(W)^\delta \subseteq X^\omega \setminus (X^* \setminus \alpha_{\mathcal{L}_\infty}(W))^\delta \subseteq \alpha_{\mathcal{L}_\infty}(W)^\delta = \mathcal{C}(W^\delta)$, and according to Property 1 the set $X^\omega \setminus (X^* \setminus \alpha_{\mathcal{L}_\infty}(W))^\delta$ is closed.

The second assertion is obvious. \square

Example 4 To see that the topology generated by $\alpha_{\mathcal{L}_\infty}$ is not strongly compatible with the CANTOR topology we remark that the closed set $\{\text{ab}\}^*$ is no (σ, δ) -subset of X^* .

Indeed, $(\{\text{ab}\}^*)^\delta$ is finite implies $\alpha_{\mathcal{L}_\infty}(\{\text{ab}\}^*) = \{\text{ab}\}^*$. \checkmark

Finally, we give a non-trivial example⁴ of an \mathcal{L} -topology not compatible with the CANTOR topology.

Example 5 Let $\mathcal{L}_{\text{CB}} := \{W : W^\delta \text{ is uncountable}\}$. As in the case of \mathcal{L}_∞ Proposition 14 and Theorem 16 prove that $\text{Anf}_{\mathcal{L}_{\text{CB}}}$ is a derived set operator. Here $(\text{Anf}_{\mathcal{L}_{\text{CB}}}(W))^\delta = \{\xi : \forall w (w \sqsubset \xi \rightarrow w \cdot X^\omega \cap W^\delta \text{ is uncountable})\}$ is the set of condensation points of W^δ in CANTOR-space (cf. [Kur66, Ch. 2, § 23, III]).

As $\text{Anf}_{\mathcal{L}_{\text{CB}}}(a^*ba^*) = \emptyset$ the language a^*ba^* is closed, but $(a^*ba^*)^\delta = a^*ba^\omega$ is not closed in CANTOR space. \checkmark

References

- [BN80] Luc Boasson and Maurice Nivat. Adherences of languages. *J. Comput. Syst. Sci.*, 20(3):285–309, 1980.
- [BN81] Luc Boasson and Maurice Nivat. Centers of languages. In Peter Deussen, editor, *Theoretical Computer Science*, volume 104 of

⁴The discrete topology $\mathcal{D} = (X^*, 2^{X^*})$ is a trivial example.

- Lecture Notes in Computer Science*, pages 245–251, Heidelberg, 1981. Springer.
- [CJS09] Cristian S. Calude, Helmut Jürgensen, and Ludwig Staiger. Topology on words. *Theoret. Comput. Sci.*, 410(24-25):2323–2335, 2009.
- [Dav64] Morton Davis. Infinite games of perfect information. In *Advances in game theory*, Edited by M. Dresher, L. S. Shapley, and A. W. Tucker. Annals of Mathematics Studies, No. 52, pages 85–101. Princeton Univ. Press, Princeton, N.J., 1964.
- [Kur66] Kazimierz Kuratowski. *Topology I*. Academic Press, New York, 1966.
- [PP04] Dominique Perrin and Jean-Éric Pin. *Infinite Words*, volume 141 of *Pure and Applied Mathematics*. Elsevier, Amsterdam, 2004.
- [Pro80] Helmut Prodinger. Topologies on free monoids induced by closure operators of a special type. *RAIRO Inform. Théor. Appl.*, 14(2):225–237, 1980.
- [Pro83] Helmut Prodinger. Topologies on free monoids induced by families of languages. *RAIRO Inform. Théor. Appl.*, 17(3):285–290, 1983.
- [PU79] Helmut Prodinger and Friedrich J. Urbanek. Language operators related to Init. *Theor. Comput. Sci.*, 8:161–175, 1979.
- [Red86] Roman R. Redziejowski. Infinite-word languages and continuous mappings. *Theor. Comput. Sci.*, 43:59–79, 1986.
- [RS63] Helena Rasiowa and Roman Sikorski. *The Mathematics of Metamathematics*. PWN, Warszawa, 1963.
- [RS08] Tobias Richter and Ludwig Staiger. Topological language operators. In *Proceedings 18. Theorietag Automaten und Formale Sprachen*, pages 109 – 114. Institut für Informatik, Universität Gießen, Gießen, 2008. <https://d-nb.info/1064837174/34#page=111>.

- [Smy92] Michael B. Smyth. Topology. In Samson Abramsky, Dov M. Gabbay, and Thomas S. E. Maibaum, editors, *Handbook of Logic in Computer Science*, volume 1, pages 641–761. Oxford Univ. Press, New York, 1992.
- [SN86] Ludwig Staiger and Werner Nehrlich. The centers of context-sensitive languages. In Jozef Gruska, Branislav Rován, and Jiří Wiedermann, editors, *Mathematical Foundations of Computer Science, 1986 (Bratislava, 1986)*, volume 233 of *Lecture Notes in Comput. Sci.*, pages 594–601. Springer, Berlin, 1986.
- [Sta87] Ludwig Staiger. Sequential mappings of ω -languages. *RAIRO Inform. Théor. Appl.*, 21(2):147–173, 1987.
- [Sta97] Ludwig Staiger. ω -languages. In Grzegorz Rozenberg and Arto Salomaa, editors, *Handbook of Formal Languages*, volume 3, pages 339–387. Springer, Berlin, 1997.
- [Sta10] Ludwig Staiger. Joint topologies for finite and infinite words. In Yuan Gao, Hanlin Lu, Shinnosuke Seki, and Sheng Yu, editors, *Developments in Language Theory*, volume 6224 of *Lecture Notes in Computer Science*, pages 442–443. Springer, 2010.
- [Tho90] Wolfgang Thomas. Automata on infinite objects. In Jan van Leeuwen, editor, *Handbook of Theoretical Computer Science*, volume B, pages 133–192. North Holland, Amsterdam, 1990.